

Ivermectin for COVID-19: real-time meta analysis of 65 studies

Covid Analysis, Oct 4, 2021, Version 127 — added Mayer (V1 Nov 26, 2020) [GMK response, Elgazzar]
<https://ivmmeta.com/>

- Meta analysis using the most serious outcome reported shows 66% [53-76%] and 86% [75-92%] improvement for early treatment and prophylaxis, with similar results after exclusion based sensitivity analysis and restriction to peer-reviewed studies or Randomized Controlled Trials.
- Statistically significant improvements are seen for mortality, ventilation, ICU admission, hospitalization, recovery, cases, and viral clearance. 30 studies show statistically significant improvements in isolation.
- Results are very robust — in worst case exclusion sensitivity analysis 54 of 65 studies must be excluded to avoid finding statistically significant efficacy.
- While many treatments have some level of efficacy, they do not replace vaccines and other measures to avoid infection. Only 26% of ivermectin studies show zero events in the treatment arm.
- Multiple treatments are typically used in combination, and other treatments could be more effective, including monoclonal antibodies which may be available in countries not recommending ivermectin (sotrovimab, casirivimab/imdevimab, and bamlanivimab/etesevimab).
- Elimination of COVID-19 is a race against viral evolution. No treatment, vaccine, or intervention is 100% available and effective for all variants. All practical, effective, and safe means should be used, including treatments, as supported by Pfizer [*Pfizer, TrialSiteNews*]. Denying the efficacy of treatments increases the risk of COVID-19 becoming endemic; and increases mortality, morbidity, and collateral damage.


	<i>Studies</i>	<i><u>Prophylaxis</u></i>	<i><u>Early treatment</u></i>	<i><u>Late treatment</u></i>	<i>Patients</i>
<u>All studies</u>	65	86% [75-92%]	66% [53-76%]	40% [24-52%]	47,741
<u>Peer-reviewed</u>	45	86% [74-93%]	70% [52-81%]	43% [21-59%]	17,316
<u>Randomized Controlled Trials</u>	32	84% [25-96%]	63% [45-75%]	30% [2-50%]	6,648

Percentage improvement with ivermectin treatment

- There is evidence of a negative publication bias, and the probability that an ineffective treatment generated results as positive as the 65 studies is estimated to be 1 in 403 billion.
- Over 20 countries have adopted ivermectin for COVID-19. The evidence base is much larger and has much lower conflict of interest than typically used to approve drugs.
- All data to reproduce this paper and sources are in the appendix. See [*Bryant, Hariyanto, Kory, Lawrie, Nardelli*] for other meta analyses with similar results confirming efficacy.



Ivermectin meta analysis mortality results

ivmmeta.com Oct 6, 2021



Evidence base used for other COVID-19 approvals			
Medication	Studies	Patients	Improvement
<u>Budesonide (UK)</u>	1	1,779	17%
<u>Remdesivir (USA EUA)</u>	1	1,063	31%
<u>Casirivimab/i.. (USA EUA)</u>	1	799	66%
<i>Ivermectin evidence</i>	65	47,717	66% [58-73%]

ivmmeta.com Oct 6, 2021






Figure 1. A. Random effects meta-analysis excluding late treatment. This plot shows pooled effects, analysis for individual outcomes is below, and more details on pooled effects can be found in the heterogeneity section. Effect extraction is pre-specified, using the most serious outcome reported. Simplified dosages are shown for comparison, these are the total dose in the first four days for treatment, and the monthly dose for prophylaxis, for a 70kg person. For details of effect extraction and full dosage information see the [appendix](#). **B.** Scatter plot showing the distribution of effects reported in early treatment studies and in all studies. **C and D.** Chronological history of all reported effects, with the probability that the observed frequency of positive results occurred due to random chance from an ineffective treatment.

Introduction

We analyze all significant studies concerning the use of ivermectin for COVID-19. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We present random effects meta-analysis results for all studies, for studies within each treatment stage, for mortality results, for COVID-19 case results, for viral clearance results, for peer-reviewed studies, for Randomized Controlled Trials (RCTs), and after exclusions.

We also perform a simple analysis of the distribution of study effects. If treatment was not effective, the observed effects would be randomly distributed (or more likely to be negative if treatment is harmful). We can compute the probability that the observed percentage of positive results (or higher) could occur due to chance with an ineffective treatment (the probability of $\geq k$ heads in n coin tosses, or the one-sided sign test / binomial test). Analysis of publication bias is important and adjustments may be needed if there is a bias toward publishing positive results.

Figure 2 shows stages of possible treatment for COVID-19. **Prophylaxis** refers to regularly taking medication before becoming sick, in order to prevent or minimize infection. **Early Treatment** refers to treatment immediately or soon after symptoms appear, while **Late Treatment** refers to more delayed treatment.




Figure 2. Treatment stages.

Results

Figure 3, 4, and 5 show results by treatment stage. Figure 6, 7, 8, 9, 10, 11, 12, and 13 show forest plots for a random effects meta-analysis of all studies with pooled effects, and for studies reporting mortality results, ICU admission, mechanical ventilation, hospitalization, recovery, COVID-19 cases, and viral clearance results only. Figure 14 shows results for peer reviewed trials only and Figure 15 shows results restricted to serious outcomes. Table 1 summarizes the results.

Treatment time	Number of studies reporting positive effects	Total number of studies	Percentage of studies reporting positive effects	Probability of an equal or greater percentage of positive results from an ineffective treatment	Random effects meta-analysis results
Early treatment	25	29	86.2%	1 in 19 thousand	66% improvement RR 0.34 [0.24-0.47] p < 0.0001
Late treatment	20	22	90.9%	1 in 17 thousand	40% improvement RR 0.60 [0.48-0.76] p < 0.0001
Prophylaxis	14	14	100%	1 in 16 thousand	86% improvement RR 0.14 [0.08-0.25] p < 0.0001
All studies	59	65	90.8%	1 in 403 billion	66% improvement RR 0.34 [0.27-0.42] p < 0.0001

Table 1. Results by treatment stage.




Figure 3. Results by treatment stage.




Figure 4. Chronological history of early and late treatment results, with the probability that the observed frequency of positive results occurred due to random chance from an ineffective treatment.







Figure 5. Chronological history of prophylaxis results.

All 65 ivermectin COVID-19 studies

ivmmeta.com Oct 6, 2021






Figure 6. Random effects meta-analysis for all studies. This plot shows pooled effects, analysis for individual outcomes is below, and more details on pooled effects can be found in the heterogeneity section. Effect extraction is pre-specified, using the most serious outcome reported. Simplified dosages are shown for comparison, these are the total dose in the first four days for treatment, and the monthly dose for prophylaxis, for a 70kg person. For details of effect extraction and full dosage information see the [appendix](#).




Figure 7. Random effects meta-analysis for mortality results only.




Figure 8. Random effects meta-analysis for mechanical ventilation results only.




Figure 9. Random effects meta-analysis for ICU admission results only.




Figure 10. Random effects meta-analysis for hospitalization results only.




Figure 11. Random effects meta-analysis for recovery results only.




Figure 12. Random effects meta-analysis for COVID-19 case results only.

All 22 ivermectin COVID-19 viral clearance results

ivmmeta.com Oct 6, 2021




Figure 13. Random effects meta-analysis for viral clearance results only.



Figure 14. Random effects meta-analysis for peer reviewed trials only. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

Randomized Controlled Trials (RCTs)

Results restricted to Randomized Controlled Trials (RCTs) are shown in Figure 16, 17, 18, and 19, and Table 2. RCT results are similar to non-RCT results. Evidence shows that non-RCT trials can also provide reliable results. [Concato] find that well-designed observational studies do not systematically overestimate the magnitude of the effects of treatment compared to RCTs. [Anglemyer] summarized reviews comparing RCTs to observational studies and found little evidence for significant differences in effect estimates. [Lee] shows that only 14% of the guidelines of the Infectious Diseases Society of America were based on RCTs. Evaluation of studies relies on an understanding of the study and potential biases. Limitations in an RCT can outweigh the benefits, for example excessive dosages, excessive treatment delays, or Internet survey bias could have a greater effect on results. Ethical issues may also prevent running RCTs for known effective treatments. For more on issues with RCTs see [Deaton, Nichol].




Figure 16. Randomized Controlled Trials. The distribution of results for RCTs is similar to the distribution for all other studies.



Figure 17. Random effects meta-analysis for Randomized Controlled Trials only. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.





Figure 19. RCTs excluding late treatment.




Figure 18. Random effects meta-analysis for Randomized Controlled Trial mortality results only.

Treatment time	Number of studies reporting positive effects	Total number of studies	Percentage of studies reporting positive effects	Probability of an equal or greater percentage of positive results from an ineffective treatment	Random effects meta-analysis results
Randomized Controlled Trials	27	32	84.4%	1 in 18 thousand	58% improvement RR 0.42 [0.29-0.59] p < 0.0001
Randomized Controlled Trials (excluding late treatment)	18	21	85.7%	1 in 1 thousand	69% improvement RR 0.31 [0.19-0.49] p < 0.0001

Table 2. Summary of RCT results.

Exclusions

To avoid bias in the selection of studies, we include all studies in the main analysis. Here we show the results after excluding studies with critical issues likely to alter results, non-standard studies, and studies where very minimal detail is currently available. Our bias evaluation is based on full analysis of each study and identifying when there is a significant chance that limitations will substantially change the outcome of the study. We believe this can be more valuable than checklist-based approaches such as Cochrane GRADE, which may underemphasize serious issues not captured in the checklists, and overemphasize issues unlikely to alter outcomes in specific cases (for example, lack of blinding for an objective mortality outcome, or certain specifics of randomization with a very large effect size). However, these approaches can be very high quality when well done, especially when the authors carefully review each study in detail [Bryant].

[Soto-Becerra] is a database analysis covering anyone with ICD-10 COVID-19 codes, which includes asymptomatic PCR+ patients. Therefore many patients in the control group are likely asymptomatic with regards to SARS-CoV-2, but in the hospital for another reason. For those that had symptomatic COVID-19, there is also likely significant confounding by indication. KM curves show that the treatment groups were in more serious condition, with more than the total excess mortality at 30 days occurring on day 1. All treatments are worse than the control group at 30 days, while at the latest followup all treatments show lower mortality than control. The machine learning system used also appears over-parameterized and likely to result in significant overfitting and inaccurate results. There is also no real control group in this study - patients receiving the treatments after 48 hours were put in the control group. Authors also state that outcomes within 24 hours were excluded, however the KM curves show significant mortality at day 1 (only for the treatment groups). Several protocol violations have also been reported in this study [Yim]. Note that this study provides both 30 day mortality and weighted KM curves up to day 43 for ivermectin, we use the day 43 results as per our protocol.

[López-Medina] has many issues. The primary outcome was changed mid-trial from clinical deterioration to complete resolution of symptoms including "not hospitalized and no limitation of activities" as a negative outcome. Critically, temporary side effects of a successful treatment may be considered as a negative outcome, which could result in falsely concluding that the treatment is not effective. Such an outcome is also not very meaningful in terms of assessing how treatment affects the incidence of serious outcomes. With the low risk patient population in this study, there is

also little room for improvement - 58% recovered within the first 2 days to "not hospitalized and no limitation of activities" or better. There was only one death (in the control arm). This study also gave ivermectin to the control arm for 38 patients and it is unknown if the full extent of the error was identified, or if there were additional undiscovered errors. The side effect data reported in this trial raises major concerns, with more side effects reported in the placebo arm, suggesting that more placebo patients may have received treatment. Ivermectin was widely used in the population and available OTC at the time of the study. The study protocol allows other treatments but does not report on usage. The name of the study drug was concealed by referring to it as "D11AX22". The presentation of this study also appears to be significantly biased. While all outcomes show a benefit for ivermectin, the abstract fails to mention that much larger benefits are seen for serious outcomes, including the original primary outcome, and that the reason for not reaching statistical significance is the low number of events in a low risk population where most recover quickly without treatment.

[Shahbaznejad] had only one death that occurred in a patient that was critically ill at the time of admission and died within the first 24 hours.

[Vallejos] reports prophylaxis results, however only very minimal details are currently available in a news report. We include these results for additional confirmation of the efficacy observed in other trials, however this study is excluded here. **[Hellwig]** analyze African countries and COVID-19 cases in October 2020 as a function of whether widespread prophylactic use of ivermectin is used for parasitic infections. **[Tanioka]** perform a similar analysis for COVID-19 mortality in January 2021. These studies are excluded because they are not clinical trials. **[Galan]** perform an RCT comparing ivermectin and other treatments with very late stage severe condition hospitalized patients, not showing significant differences between the treatments. Authors were unable to add a control arm due to ethical issues. The closest control comparison we could find is **[Baqui]**, which shows 43% hospital mortality in the northern region of Brazil where the study was performed, from which we can estimate the mortality with ivermectin in this study as 47% lower, RR 0.53. Further, the study is restricted to more severe cases, hence the expected mortality, and therefore the benefit of treatment, may be higher. **[Kishoria]** restrict inclusion to patients that did not respond to standard treatment, provide no details on the time of the discharge status, and there are very large unadjusted differences in the groups, with over twice as many patients in the ivermectin group with age >40, and all patients over 60 in the ivermectin group.

Summarizing, the studies excluded are as follows, and the resulting forest plot is shown in Figure 20. [The supplementary data](#) contains results after restrictions and exclusions.

[Ahsan], unadjusted results with no group details.

[Cadegiani], control group retrospectively obtained from untreated patients in the same population.

[Carvalho], concern about potential data issues.

[Carvalho (B)], concern about potential data issues.

[Carvalho (C)], minimal details of groups provided.

[Elavarasi], unadjusted results with no group details.

[Hazan], study uses a synthetic control arm.

[Hellwig], not a typical trial, analysis of African countries that used or did not use ivermectin prophylaxis for parasitic infections.

[Kishoria], excessive unadjusted differences between groups.

[López-Medina], strong evidence of patients in the control group self-medicating, ivermectin widely used in the population at that time, and the study drug identity was concealed by using the name D11AX22.

[Roy], no serious outcomes reported and fast recovery in treatment and control groups, there is little room for a treatment to improve results.

[Soto-Becerra], substantial unadjusted confounding by indication likely, includes PCR+ patients that may be asymptomatic for COVID-19 but in hospital for other reasons.

[Szente Fonseca], result is likely affected by collinearity across treatments in the model.

[Tanioka], not a typical trial, analysis of African countries that used or did not use ivermectin prophylaxis for parasitic infections.

[Together Trial], preliminary report with minimal details.

[Vallejos], detail too minimal.

49 ivermectin COVID-19 studies after exclusions

ivmmeta.com Oct 6, 2021




Figure 20. Random effects meta-analysis excluding studies with significant issues. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

Heterogeneity

Heterogeneity in COVID-19 studies arises from many factors including:

Treatment delay. The time between infection or the onset of symptoms and treatment may critically affect how well a treatment works. For example an antiviral may be very effective when used early but may not be effective in late stage disease, and may even be harmful. Figure 21 shows an example where efficacy declines as a function of treatment delay. Other medications might be beneficial for late stage complications, while early use may not be effective or may even be harmful. Oseltamivir, for example, is generally only considered effective for influenza when used within 0-36 or 0-48 hours [McLean, Treanor].




Figure 21. Effectiveness may depend critically on treatment delay.

Patient demographics. Details of the patient population including age and comorbidities may critically affect how well a treatment works. For example, many COVID-19 studies with relatively young low-comorbidity patients show all patients recovering quickly with or without treatment. In such cases, there is little room for an effective treatment to improve results (as in [López-Medina]).

Effect measured. Efficacy may differ significantly depending on the effect measured, for example a treatment may be very effective at reducing mortality, but less effective at minimizing cases or hospitalization. Or a treatment may have no effect on viral clearance while still being effective at reducing mortality.

Variants. There are thousands of different variants of SARS-CoV-2 and efficacy may depend critically on the distribution of variants encountered by the patients in a study.

Regimen. Effectiveness may depend strongly on the dosage and treatment regimen. Higher dosages have been found to be more successful for ivermectin [Babalola]. Method of administration may also be critical. [Guzzo] show that the plasma concentration of ivermectin is much higher when administered with food (Figure 22: geometric mean AUC 2.6 times higher). Many ivermectin studies specify fasting, or they do not specify administration. Fasting administration is expected to reduce effectiveness for COVID-19 due to lower plasma and tissue concentrations. Note that this is different to anthelmintic use in the gastrointestinal tract where fasting is recommended.

Treatments. The use of other treatments may significantly affect outcomes, including anything from supplements, other medications, or other kinds of treatment such as prone positioning.




Figure 22. Mean plasma concentration (ng/mL) profiles of ivermectin following single oral doses of 30mg (fed and fasted administration), from [Guzzo].

The distribution of studies will alter the outcome of a meta analysis. Consider a simplified example where everything is equal except for the treatment delay, and effectiveness decreases to zero or below with increasing delay. If there are many studies using very late treatment, the outcome may be negative, even though the treatment may be very effective when used earlier.

In general, by combining heterogeneous studies, as all meta analyses do, we run the risk of obscuring an effect by including studies where the treatment is less effective, not effective, or harmful.

When including studies where a treatment is less effective we expect the estimated effect size to be lower than that for the optimal case. We do not *a priori* expect that pooling all studies will create a positive result for an effective treatment. Looking at all studies is valuable for providing an overview of all research, and important to avoid cherry-picking, but the resulting estimate does not apply to specific cases such as early treatment in high-risk populations.

Ivermectin studies vary widely in all the factors above, which makes the consistently positive results even more remarkable. A failure to detect an association after combining heterogeneous studies does not mean the treatment is not effective (it may only work in certain cases), however the reverse is not true – an identified association is valid, although the magnitude of the effect may be larger for more optimal cases, and lower for less optimal cases. As above, the probability that an ineffective treatment generated results as positive as the 65 studies to date is estimated to be 1 in 403 billion. This result benefits from the fact that ivermectin shows some degree of efficacy for

COVID-19 in a wide variety of cases. It also likely benefits from the fact that relatively few ivermectin trials to date have been designed in a way that favors poor results. However, more trials designed in this way are expected, for example the TOGETHER trial is testing ivermectin in locations known to have a high degree of self-medication and using low doses compared to current clinical recommendations as updated for current variants. As with a companion trial, this trial may also include very low-risk patients, include relatively late treatment while identifying as an early treatment trial, and use an active placebo (vitamin C). While we present results for all studies in this paper, the individual outcome and treatment time analyses are more relevant for specific use cases.

Discussion

Publication bias. Publishing is often biased towards positive results, which we would need to adjust for when analyzing the percentage of positive results. For ivermectin, there is currently not enough data to evaluate publication bias with high confidence. One method to evaluate bias is to compare prospective vs. retrospective studies. Prospective studies are likely to be published regardless of the result, while retrospective studies are more likely to exhibit bias. For example, researchers may perform preliminary analysis with minimal effort and the results may influence their decision to continue. Retrospective studies also provide more opportunities for the specifics of data extraction and adjustments to influence results. Figure 23 shows a scatter plot of results for prospective and retrospective studies. The median effect size for prospective studies is 74% improvement, compared to 73% for retrospective studies, showing no significant difference. *[Bryant]* also perform a funnel plot analysis, which they found did not suggest evidence of publication bias. We note that there is extreme attention from many people towards locating any missing or unpublished negative trials — if they existed it would likely be known. Negative studies are submitted to us by multiple people immediately on publication. On the other hand, there is substantial evidence that journals are delaying the publication of positive studies, for example by accepting a paper for review, holding it for some time, and then rejecting it without review *[Jerusalem Post, Kory (B)]*.




Figure 23. Prospective vs. retrospective studies.

News coverage of ivermectin studies is extremely biased. Only one study to date has received significant press coverage in western media *[López-Medina]*, which is neither the largest or the least biased study, and is one of the two studies with the most critical issues as discussed earlier.

In Vitro evidence on required concentration. Some people claim that *[Caly]* shows that therapeutic concentrations are not easily reached in humans. This is incorrect. The authors explain why their *in vitro* study cannot be used to determine the effective dose *in vivo*, and state that the concentration required is very unlikely to be an issue *[Wagstaff]*. The study used monkey kidney cells (the only choice at the time of the experiments), which lack adaptive immune responses and do not produce interferon. Authors also note that ivermectin accumulates in lung and other tissues, that

subsequent experiments with lung cells show many times greater concentrations, and that the average lung concentration shown in modeling studies exceeds the effective level shown in their research. Authors note that ivermectin works with the immune system and a 1:1 ratio of drug to virus is unlikely to be required. In *[Bray]*, author reply that "ivermectin's key direct target in mammalian cells is a not a viral component, but a host protein important in intracellular transport; the fact that it is a host-directed agent (HDA) is almost certainly the basis of its broad-spectrum activity against a number of different RNA viruses in vitro. The way a HDA can reduce viral load is by inhibiting a key cellular process that the virus hijacks to enhance infection by suppressing the host antiviral response. Reducing viral load by even a modest amount by using a HDA at low dose early in infection can be the key to enabling the body's immune system to begin to mount the full antiviral response before the infection takes control." In further research, authors note that they find efficacy for prophylactic use, and that smaller repeated doses are more effective than a single larger dose *[Wagstaff]*.

Conflicts of interest. Pharmaceutical drug trials often have conflicts of interest whereby sponsors or trial staff have a financial interest in the outcome being positive. Ivermectin for COVID-19 lacks this because it is off-patent, has many manufacturers, and is very low cost. In contrast, most COVID-19 ivermectin trials have been run by physicians on the front lines with the primary interest of finding the best methods to save human lives and minimize the collateral damage caused by COVID-19. While pharmaceutical companies are careful to run trials under optimal conditions (for example, restricting patients to those most likely to benefit, only including patients that can be treated soon after onset when necessary, ensuring accurate dosing), many ivermectin trials do not represent the optimal conditions for efficacy.

Two ivermectin trials to date involve very large financial conflicts of interest *[López-Medina, Together Trial]* — companies closely involved with the trial or organizers stand to lose billions of dollars if ivermectin efficacy becomes more widely known. The design of these trials favors producing a null outcome as detailed in *[López-Medina, Together Trial]*. Note that biasing an RCT to produce a false positive result is difficult (suppressing adverse events is relatively easy *[Evans]*), but biasing a trial to produce a false negative result is very easy — for example, in a trial of an antiviral that works within the first 24 hours of symptom onset, trial organizers only need to avoid treating people within the first 24 hours; or with a disease like COVID-19, organizers only need to select a low-risk population where most people recover quickly without treatment. We note that, even under the very suboptimal designs, these trials produced positive results, although without statistical significance.

Early/late vs. mild/moderate/severe. Some analyses classify treatment based on early/late administration (as we do here), while others distinguish between mild/moderate/severe cases. We note that viral load does not indicate degree of symptoms — for example patients may have a high viral load while being asymptomatic. With regard to treatments that have antiviral properties, timing of treatment is critical — late administration may be less helpful regardless of severity.

Study notes. 3 of the 65 studies compare against other treatments rather than placebo. Currently ivermectin shows better results than these other treatments, however ivermectin may show greater improvement when compared to placebo. 15 of 65 studies combine treatments, for example ivermectin + doxycycline. The results of ivermectin alone may differ. 4 of 32 RCTs use combined treatment, three with doxycycline, and one with iota-carrageenan. 1 of 65 studies currently have minimal published details available.

Meta analyses. Typical meta analyses involve subjective selection criteria, effect extraction rules, and study bias evaluation, which can be used to bias results towards a specific outcome. In order to avoid bias we include all studies and use a pre-specified method to extract results from all studies (we also present results after exclusions). The results to date are overwhelmingly positive, very consistent, and very insensitive to potential selection criteria, effect extraction rules, and/or bias evaluation. Additional meta analyses confirming the effectiveness of ivermectin can be found in [Bryant, Kory, Lawrie]. Figure 24 shows a comparison of mortality results across meta analyses. [Kory] also review epidemiological data and provide suggested treatment regimens.




Figure 24. Comparison of mortality results from different meta analyses. OR converted to RR for [Kory, Nardelli]. OR displayed for [WHO]. WHO provides two results, one based on 5 studies and one based on 7, with no explanation for the difference. The result based on 7 studies is shown here, for which the details required to calculate the RR are not provided.

Evidence base. The evidence supporting ivermectin for COVID-19 far exceeds the typical amount of evidence used for the approval of treatments. [Lee] shows that only 14% of the guidelines of the Infectious Diseases Society of America were based on RCTs. Table 3 and Table 4 compare the amount of evidence for ivermectin compared to that used for other COVID-19 approvals, and that used by WHO for the approval of ivermectin for scabies and strongyloidiasis. Table 5 compares US CDC recommendations for ibuprofen and ivermectin.

Indication	Studies	Patients	Status
Strongyloidiasis [Kory (C)]	5	591	Approved
Scabies [Kory (C)]	10	852	Approved
COVID-19	65	47,717	Pending
COVID-19 RCTs	32	6,648	

Table 3. WHO ivermectin approval status.

Medication	Studies	Patients	Improvement	Status
<u>Budesonide (UK)</u>	1	1,779	17%	Approved
<u>Remdesivir (USA)</u>	1	1,063	31%	Approved
<u>Casiri/imdevimab (USA)</u>	1	799	66%	Approved
<i>Ivermectin evidence</i>	65	47,717	66% [58-73%]	Pending

Table 4. Evidence base used for other COVID-19 approvals compared with the ivermectin evidence base.

	<i>Ibuprofen</i>	<i>Ivermectin (for scabies)</i>	<i>Ivermectin (for COVID-19)</i>
Lives saved	0	0	>500,000
Deaths per year	~450	<1	<1
CDC recommended	Yes	Yes	No
Based on	0 RCTs	10 RCTs 852 patients	32 RCTs 6,648 patients

Table 5. Comparison of CDC recommendations [Kory (C)].

WHO Analysis

WHO updated their treatment recommendations on 3/30/2021 [WHO]. For ivermectin they reported a mortality odds ratio of 0.19 [0.09-0.36] based on 7 studies with 1,419 patients. They do not specify which trials they included. The report is inconsistent, with a forest plot that only shows 4 studies with mortality results. WHO's recommendation has not been updated for 190 days.

Despite this extremely positive result, they recommended only using ivermectin in clinical trials. The analysis contains many flaws [Kory (D)]:

- Of the 65 studies (32 RCTs), they only included 16.
- They excluded all 14 prophylaxis studies (3 RCTs).
- There was no protocol for data exclusion.
- Trials included in the original UNITAID search protocol were excluded.
- They excluded all epidemiological evidence, although WHO has considered such evidence in the past.
- They combine early treatment and late treatment studies and do not provide heterogeneity information. As above, early treatment is more successful, so pooling late treatment studies will obscure the effectiveness of early treatment. They chose not to do subgroup analysis by disease

severity across trials, although treatment delay is clearly a critical factor in COVID-19 treatment, the analysis is easily done (as above), and it is well known that the studies for ivermectin and many other treatments clearly show greater effectiveness for early treatment.

- WHO downgraded the quality of trials compared to the UNITAID systematic review team and a separate international expert guideline group that has long worked with the WHO **[Bryant]**.
- They disregarded their own guidelines that stipulate quality assessments should be upgraded when there is evidence of a large magnitude effect (which there is), and when there is evidence of a dose-response relationship (which there is). They claim there is no dose-response relationship, while the UNITAID systematic review team found a clear relationship, along with individual studies **[Babalola]**.
- Their risk of bias assessments do not match the actual risk of bias in studies. For example they classify **[López-Medina]** as low risk of bias, however this study has many issues making the results unreliable **[Covid Analysis]**, even prompting an open letter from over 170 physicians concluding that the study is fatally flawed **[Open Letter]**. **[Gonzalez]** is also classified as low risk of bias, but is a study with very late stage severe condition high-comorbidity patients. There is a clear treatment delay-response relationship and very late stage treatment is not expected to be as effective as early treatment. Conversely, much higher quality studies were classified as high risk of bias.
- Although WHO's analysis is called a "living guideline", it is rarely updated and very out of date. As of May 14, 2021, four of the missing RCTs are known to WHO and labeled "RCTs pending data extraction" **[COVID-NMA]**. We added these 4, 4, 2, and one month earlier.
- A single person served as Methods Chair, member of the Guidance Support Collaboraton Committee, and member of the Living Systematic Review/NMA team.
- Public statements from people involved in the analysis suggest substantial bias. For example, a co-chair reportedly said that "the data available was sparse and likely based on chance" **[Reuters]**. As above, the data is comprehensive, and we estimate the probability that an ineffective treatment generated results as positive as observed to be 1 in 403 billion. The clinical team lead refers to their analysis of ivermectin as "fighting this overuse of unproven therapies ... without evidence of efficacy" **[Reuters]**, despite the extensive evidence of efficacy from the 65 studies by 655 scientists with 47,717 patients. People involved may be more favorable to late stage treatment of COVID-19, for example the co-chair recommended treating severe COVID-19 with remdesivir **[Rochwerg]**.

In summary, although WHO's analysis predicts that over 2 million fewer people would be dead if ivermectin was used from early in the pandemic, they recommend against use outside trials. This appears to be based primarily on excluding the majority of the evidence, and by assigning bias estimates that do not match the actual risk of bias in studies.

Use early in the pandemic was proposed by Kitasato University including the co-discoverer of ivermectin, Dr. Satoshi Ōmura. They requested Merck conduct clinical trials of ivermectin for COVID-19 in Japan, because Merck has priority to submit an application for an expansion of ivermectin's indications. Merck declined **[Yagisawa]**.

Merck Analysis

Merck has recommended against ivermectin [Merck], however this recommendation has not been updated for 244 days.

They stated that there is "no scientific basis for a potential therapeutic effect against COVID-19 from pre-clinical studies". This is contradicted by many papers and studies, including [Arévalo, Bello, Choudhury, de Melo, DiNicolantonio, DiNicolantonio (B), Errecalde, Eweas, Francés-Monerris, Heidary, Jans, Jeffreys, Kalfas, Kory, Lehrer, Li, Mody, Mountain Valley MD, Qureshi, Saha, Surnar, Udofia, Wehbe, Yesilbag, Zaidi, Zatloukal].

They state that there is "no meaningful evidence for clinical activity or clinical efficacy in patients with COVID-19 disease". This is contradicted by numerous studies including [Afsar, Alam, Aref, Babalola, Behera, Behera (B), Bernigaud, Budhiraja, Bukhari, Chaccour, Chahla, Chahla (B), Chowdhury, Elalfy, Espitia-Hernandez, Faisal, Hashim, Huvemek, Khan, Lima-Morales, Loue, Mahmud, Mayer, Merino, Mohan, Mondal, Morgenstern, Mourya, Niaee, Okumuş, Ravikirti, Samaha, Seet].

They also claim that there is "a concerning lack of safety data in the majority of studies". Safety analysis is found in [Descotes, Errecalde, Guzzo, Kory, Madrid], and safety data can be found in most studies, including [Abd-El salam, Afsar, Ahmed, Aref, Babalola, Behera (B), Bhattacharya, Biber, Bukhari, Camprubí, Carvallo (C), Chaccour, Chahla (B), Chowdhury, Elalfy, Espitia-Hernandez, Gorial, Hazan, Huvemek, Khan, Kishoria, Krolewiecki, Lima-Morales, Loue, López-Medina, Mahmud, Mohan, Morgenstern, Mourya, Niaee, Okumuş, Pott-Junior, Seet, Shahbaznejad, Shouman, Spoorthi, Szenté Fonseca, Vallejos (B)].

Merck has a number of conflicts of interest:

- Merck has committed to give ivermectin away for free "as much as needed, for as long as needed" in the Mectizan® Donation Program [Merck (B)], to help eliminate river blindness.
- Merck has their own new COVID-19 treatments MK-7110 (formerly CD24Fc) [Adams] and Molnupiravir (MK-4482) [Merck (C), Wikipedia]. Merck has a ~\$US1.2B agreement to supply molnupiravir to the US government, if it receives EUA or approval [Khan (B)]. Over \$US10B in near-term orders are expected if approved [Genetic Engineering and Biotechnology News].
- Ivermectin is off-patent, there are many manufacturers, and Merck is unlikely to be able to compete with low cost manufacturers.
- Promoting the use of low cost off-patent medications compared to new products may be undesirable to some shareholders.
- Japan requested Merck conduct clinical trials early in the pandemic and they declined. Merck may be reluctant to admit this mistake [Yagisawa].

FDA Analysis

The US FDA recommended against ivermectin on March 5, 2021, however they stated that "*The FDA has not reviewed data to support use of ivermectin in COVID-19 patients to treat or to prevent COVID-19*". There is still no indication that the FDA has reviewed the clinical trials 215 days later.

The FDA notes that they "*received multiple reports of patients who have required medical support and been hospitalized after self-medicating with ivermectin intended for horses*". The number of reports was 4 [Pfeiffer]. For comparison, acetaminophen overdose results in ~33,000 yearly hospitalizations in the USA (~12,000 unintentional) [Charilaou]. The FDA's recommendation may increase cases of self-medication with animal ivermectin, because it reduces the percentage of prescribing physicians.

They state that "*Ivermectin is not an anti-viral*", however many studies contradict this [Ahmed, Aref, Babalola, Biber, Bukhari, Buonfrate, Caly, Chowdhury, Elalfy, Espitia-Hernandez, Khan, Mahmud, Mohan, Mourya, Okumuş], including 10 RCTs.

They note that "*some initial research is underway*", however there had been many studies completed and published prior to the FDA recommendation [Afsar, Ahmed, Alam, Babalola, Behera, Bernigaud, Biber, Budhiraja, Bukhari, Cadegiani, Camprubí, Carvalho (C), Chaccour, Chachar, Chahla (B), Chowdhury, Elalfy, Espitia-Hernandez, Gonzalez, Gorial, Hashim, Hellwig, Khan, Lima-Morales, López-Medina, Mahmud, Mohan, Niaee, Okumuş, Podder, Rajter, Ravikirti, Samaha, Shouman, Spoorthi], including 19 RCTs.

Sep 3, 2021: The FDA revised their statement slightly. They removed the false claim that ivermectin is not an antiviral, and they removed the statement that they have not reviewed the data. However, there is still nothing to indicate that they have reviewed the clinical trials. Indeed, they state "currently available data do not show ivermectin is effective against COVID-19" and "ivermectin has not been shown to be safe or effective for these indications", which are both false.

Conclusion

Ivermectin is an effective treatment for COVID-19. Treatment is more effective when used early. Meta analysis using the most serious outcome reported shows 66% [53-76%] and 86% [75-92%] improvement for early treatment and prophylaxis, with similar results after exclusion based sensitivity analysis and restriction to peer-reviewed studies or Randomized Controlled Trials. Statistically significant improvements are seen for mortality, ventilation, ICU admission, hospitalization, recovery, cases, and viral clearance. 30 studies show statistically significant improvements in isolation. The consistency of positive results across a wide variety of heterogeneous studies is remarkable, with 91% of the 65 studies reporting positive effects.

Responses

Inconclusive meta analyses. [Popp, Roman] provide meta analyses that show positive effects without reaching statistical significance. The primary methods used that result in a lack of statistical significance are the exclusion of the majority of the evidence base, and division of the remaining subset. For more details see the study notes.

Meta analysis should not combine heterogeneous studies. All meta analyses combine heterogeneous studies, because all studies differ in one or more ways, including patient demographics, treatment delay distribution, effect measured, SARS-CoV-2 variants, and treatment regimens (note that this is different to heterogeneity caused by bias). Combining heterogeneous studies may obscure efficacy - for example if treatment within 24 hours is twice as effective as treatment within 48 hours and we include studies with later treatment; or if a treatment is effective at reducing mortality but has no effect on viral clearance and we include viral clearance studies. Including studies that are further from the optimal treatment situation will reduce the observed effect size. This can be seen in the treatment delay analysis - late treatment is less effective and including late treatment studies lowers the effect size. For any negative meta analysis, we must consider if the treatment is effective but only in a subset of the situations covered by the studies (or a situation not covered by any study, for example few treatments have studies with a treatment delay ≤ 24 hours).

How should the result be interpreted when pooling effects? In the pooled analysis, the result is a weighted average of the improvement in the most serious outcome reported. The specific analyses should be used for specific outcomes. Note that a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases, etc. Note that we have to consider all information to create the most accurate prediction of efficacy. While there are more sophisticated ways to combine all of the information, the advantage of the method used here is simplicity and transparency. Note that the highly significant results observed are without incorporating additional information that would further increase confidence, such as the treatment delay-response relationship.

Primary outcome analysis. We use fixed pre-specified effect extraction to avoid bias and to focus on the most clinically relevant results. For comparison, we have also performed analysis using the primary outcome of studies, with results showing similar effect sizes. Prophylaxis results are very similar with 100% (14 of 14) positive effects. Early treatment shows 97% (28 of 29) positive effects, improved due to the very small event count negative serious outcomes in Krolewiecki, Vallejos, and Buonfrate no longer having priority. Late treatment shows 77% (17 of 22) positive effects, reduced slightly, primarily due to viral clearance results being the primary outcome in some studies, and viral clearance being less successful with late treatment. Overall, the primary outcome analysis shows 91% (59 of 65) positive effects, which is currently identical to the results of the main protocol analysis.

Elgazzar. This study was withdrawn and was removed from this analysis on the same day. There was no significant change (excluding 1 of 66 studies has very little effect, and the exclusion actually improves the treatment delay-response relationship).

Carvallo. An influential anti-treatment Twitter personality has raised concerns about [Carvallo]. There appears to be some valid concerns with potential data issues. Exclusion-based analysis shows no significant change in results, with only a minor reduction in prophylaxis efficacy to 84% [69-91%]. However, it is difficult to trust information from this personality. The author suggests that the study may not have happened at all, claiming for example that the team could not have afforded the medications without funding, and that a busy clinician would not have enough time. However, with just basic checks, the author would know that a drug company has confirmed donating the medications, that they confirmed authorization for the study was received, that the main hospital for the study requested additional supplies, and that the hospital confirmed ethics committee approval. For additional details see [O'Reilly]. We also note that the combined treatment in this study has been independently shown to be effective, and the complementary mechanisms of action support improved efficacy of the combination [Figueroa].

AT response. A technology blog published an article with incorrect and unsupported claims. The article refers to c19ivermectin.com (which is only a database of ivermectin research), but makes comments about this analysis. Most of the comments in this article are already addressed above.

Author correctly notes that the majority of results are positive and that no matter how you slice the data, the results are positive, but appears to dismiss the obvious reason without examining the evidence.

Author believes that because other effective treatments exist, and because we have also covered those, there must be a positive bias. For ivermectin though, we find evidence of a negative publication bias, and despite enormous worldwide attention, there is no evidence of missing negative trials, while there is substantial evidence of positive trials being delayed by editors (journals fast track null results, while holding positive trials and later returning them without review). We also note that many of the effective treatments are adopted by governments worldwide, including several in the author's country. Approved treatments include sotrovimab, casirivimab, imdevimab, bamlanivimab, etesevimab, budesonide, favipiravir, and convalescent plasma (although not showing efficacy in our analysis), others have already been purchased pending approval or are not yet available (molnupiravir, proxalutamide), and others are widely accepted to be helpful, including in the author's country, despite gaining minimal attention from authorities (vitamin D, vitamin C) [Miller].

Author finds the heterogeneity in dosage, treatment time, etc. concerning. This heterogeneity is beneficial and gives us much more information on the situations where treatment is effective, and the optimal dosage. Results from a single study only apply to the conditions of that study and cannot be extrapolated to other conditions — author makes this mistake claiming another treatment is ineffective based on definitive evidence, but that evidence only applies to very late treatment in a very sick population with excessive dosage — not the optimal use of an antiviral for COVID-19. While we cannot use the larger evidence base to predict a specific situation, e.g., mortality in high risk patients with specific treatment delay and dosing, we can use the larger evidence base as evidence for/against efficacy, and many subgroup analyses have sufficient evidence for more specific cases.

Author refers to the withdrawn Elgazzar study (removed from this analysis on the same day) as a major development, however there was no significant change. Excluding 1 of 66 studies has very little effect, and the exclusion actually improves the treatment delay-response relationship. 54 of 65 studies need to be excluded to avoid finding statistically significant efficacy in a worst case sensitivity analysis.

Author is concerned that some studies use combined treatment, however 50 do not use combined treatment, and most of the additions are treatments independently known to not have significant efficacy alone.

We also note that the author has never contacted us.

Twitter personality response. An influential anti-treatment Twitter personality, journalist, and student epidemiologist has made a number of incorrect, misleading, hyperbolic, and unsupported statements. Author has been paid for writing anti-treatment articles and has referred to ivermectin as a "horse dewormer".

Author's attempt to discredit ivermectin research centers on the fundamentally false assertion that excluding a small number of lower quality trials results in a negative outcome. It should be clear from the forest plot that this is not possible, but we can be more specific. We perform an absolute worst case sensitivity analysis, where positive studies are excluded in order of the effect size, with

the largest effect first. How many studies do we need to exclude before the meta analysis RR has a confidence interval exceeding 1.0? **83%, or 54 of 65 studies must be excluded to avoid finding statistically significant efficacy.** As with all data in this paper, this analysis will automatically update as the evidence base evolves. Also note that this is after exclusion of the two currently withdrawn papers - one has never been in this analysis, the second was removed on the same day it was withdrawn.

Author makes a basic error by equating positive effects that are not statistically significant at a specific level with "no effect", a misunderstanding of statistics [*Amrhein*]. For example, if a study reports 50% improvement with a *p* value of 0.1, we cannot say that the study shows the treatment is ineffective, or in the words of the author shows "no benefit at all". Author repeatedly makes false claims in this way.

The author's other errors, none of which have been corrected over a month later, include:

- that excluding Elgazzar et al. completely changes the results (excluding 1 of 66 studies has very little effect, and the exclusion actually improves the treatment delay-response relationship)
- making basic errors suggesting very superficial reading of studies, for example claiming the RR in Szente Fonseca is the risk of being treated
- making basic errors suggesting very superficial reading of this paper, for example claiming that a result for prophylaxis studies is based on the number of patients from all studies
- equating a high degree of COVID-19 in a country partially adopting a treatment with a lack of efficacy, disregarding obvious confounding such as heavily affected areas being more likely to adopt treatment (analysis of results in regions or time periods adopting treatment, while not equivalent to controlled studies, is more informative and shows efficacy [*Chamie-Quintero, Chamie-Quintero (B), Merino, Ontai*])
- confusing heterogeneity due to dose, treatment delay, etc. and due to bias
- disregarding treatment delay to dilute or obscure effects by including late treatment (author has also used this method with other treatments)
- disregarding the existence of specific outcome analyses, RCT analysis, and exclusion-based sensitivity analysis
- suggesting that efficacy over longer periods is not possible because ivermectin has a half-life of "about a day". Author disregards known efficacy for other conditions over much longer periods, and mischaracterizes the half-life. Antiparasitic efficacy can persist for several months after a single dose [*Canga*]. Plasma half-life is longer in some studies, and significant plasma concentration can persist for over 2 weeks in some patients [*Muñoz*]. More importantly, ivermectin is highly lipophilic and may accumulate in the lung and other tissues [*Chaccour (B)*].
- misunderstanding funnel plot analysis and explanations other than selective reporting (and providing no evidence of unreported negative studies, while there is substantial evidence of difficulty publishing positive studies [*Jerusalem Post, Kory (B)*])
- suggesting that it is impossible to combine evidence from mortality and hospitalization (for example), but combining late treatment and early treatment in order to obscure efficacy (if a treatment reduces disease severity requiring hospitalization, reduced mortality in at-risk populations logically follows, whereas lack of efficacy several days after onset can not be extrapolated to early treatment — treatments for a viral infection are often less effective when delayed)

- making serious claims about individual studies without contacting authors (for example claiming patients were excluded for reaching the endpoint too quickly, whereas authors report exclusions due to baseline negative status)
- author is unaware of different variants, suggesting that results should be identical for treatment at a given delay, even when the predominant variants have markedly different peak viral load, time to peak viral load [*Faria, Karita, Nonaka*], and mortality (for example Gamma vs. non-Gamma aHR 4.73 [1.15-19.41] [*Zavascki*])
- suggesting that a specific symptom such as cough should be used (author would prefer a less positive result for the study)
- suggesting that viral load is more important than symptomatic results
- suggesting that mortality should be used in populations with zero mortality (for low-risk populations with no mortality, reduction in mortality is not possible, this does not mean a reduction in hospitalization, for example, is not valuable)
- suggesting that unadjusted results should be used in a study where the adjustments clearly make a significant difference (author wants to cherry-pick unadjusted cough results)
- suggesting that, for example, in a study of viral load where all patients recover, it is not valuable if treated patients recover faster (or are less likely to transmit the virus to others)
- suggesting that study selected outcomes should have priority rather than using a consistent pre-specified protocol, disregarding the added bias and the fact that this actually improves results for ivermectin (for example the very small event count negative serious outcomes in Krolewiecki, Vallejos, and Buonfrate would no longer have priority)
- suggesting that cough is a more important symptom than low SpO₂ or fever. Cough can persist for a long time after more serious symptoms resolve, and persistent cough may be caused by many conditions.
- suggesting that combined low dose treatment results should be used in a study that had a combined ivermectin/doxycycline arm (single dose ivermectin, 5 days doxycycline) and an ivermectin arm with treatment for 5 days
- suggesting that hospitalization time should be used for symptomatic recovery in a study where discharge is based on viral clearance (and only tested weekly).

We note that this personality has an extensive history of incorrect advice, including for example:

- claiming that flu is more dangerous than COVID-19
- claiming that SARS-CoV-2 is not airborne
- claiming that it's impossible to improve immune system functioning
- even believing and propagating a made up story that claimed ivermectin overdose was causing gunshot victims to wait at an ER

Author has taken a public position against early treatments for COVID-19 since at least July 2020. Given this longstanding and influential negative position, they may tend to view information with a negative filter and confirmation bias, and may be reluctant to admit errors. They acknowledge not having read all of the studies (and appear to have very superficially read others). They submitted zero feedback to us, suggesting that they know their comments are incorrect or that they have a

motivation other than correcting errors. Author claims that they could not contact us, however there are over 50 feedback links throughout this article. We also note that the author is not open to critical feedback and routinely blocks Twitter users correcting mistakes or expressing anything critical on their feed.

Based on many comments, author appears to focus on superficial criteria such as typesetting and quality of writing. While many of the studies have been performed by non-native English speakers with minimal budgets, this does not imply the researchers are less reliable. Indeed, the author is highly critical of the program used to create a graph, for example, but is unable to see flaws in high budget high conflict of interest trials, even when they prompt >100 scientists to write an open letter requesting retraction [*Open Letter*].


Study Notes

Together Trial.

Minimal information about the Together Trial is currently available. They released partial results in a presentation, but have not released the preprint yet. The preprint for the fluvoxamine arm, concluded at the same time, was released August 23, 2021.

The trial randomization chart does not match the protocol, suggesting major problems and indicating substantial confounding by time. For example, trial week 43, the first week for 3 dose ivermectin, shows ~3x patients assigned to ivermectin vs. placebo [*reddit.com*]. Treatment efficacy can vary significantly over time, for example due to overall improvement in protocols, changes in the distribution of variants, or changes in public awareness and treatment delays. [*Zavascki*] show dramatically higher mortality for Gamma vs non-Gamma variants (28 day mortality from symptom onset aHR 4.73 [1.15-19.41]), and the prevalence of the Gamma variant varied dramatically throughout the trial [*ourworldindata.org*]. This introduces confounding by time, which is common in COVID-19 retrospective studies and has often obscured efficacy (many retrospectives have more patients in the treatment group earlier in time when overall treatment protocols were significantly worse).

According to this analysis [*reddit.com*], the total number of patients for the ivermectin and placebo groups do not appear to match the totals in the presentation (the numbers for the fluvoxamine arm match) — reaching the number reported for ivermectin would require including some of the patients assigned to single dose ivermectin. Reaching the placebo number requires including placebo patients from the much earlier ivermectin single dose period, and from the early two week period when zero ivermectin patients were assigned. If these earlier participants were accidentally included in the control group, this would dramatically change the results in favor of the control group according to the changes in Gamma variant prevalence.



Treatment delay is currently unknown, however the protocol allows very late inclusion and a companion trial reported mostly late treatment. Overall mortality is high for 18+ outpatients. Results may be impacted by late treatment, poor SOC, and may be specific to local variants [Faria, Nonaka, Sabino]. Treatment was administered on an empty stomach, greatly reducing expected tissue concentration [Guzzo] and making the effective dose about 1/5th of current clinical practice. The trial was conducted in Minas Gerais, Brazil which had substantial community use of ivermectin [otempo.com.br], and prior use of ivermectin is not listed in the exclusion criteria.

Time from symptom onset to randomization is specified as within 7 days. However the schedule of study activities specifies treatment administration only one day after randomization, suggesting that treatment was delayed an additional day for all patients.

This trial uses a soft primary outcome, easily subject to bias and event inflation in both arms (e.g., observe >6 hours independent of indication). There is also an unusual inclusion criteria: "patients with expected hospital stays of ≤ 5 days". This is similar to "patients less likely to need treatment beyond SOC to recover", and would make it very easy to reduce the effect seen. This is not in either of the published protocols.

The trial took place in an area of Brazil where the Gamma variant was prominent. Brazilian clinicians report that this variant is much more virulent, and that significantly higher dosage and/or earlier treatment is required, as may be expected for variants where the peak viral load is significantly higher and/or reached earlier [Faria, Nonaka].

Trial design, analysis, and presentation, along with previous public and private statements suggest investigator bias. Design: including very late treatment, additional day before administration, operation in a region with high community use, specifying administration on an empty stomach, limiting treatment to 3 days, using soft inclusion criterion and a soft primary outcome, easily subject to bias. Analysis: authors perform analysis excluding events very shortly after randomization for fluvoxamine but not ivermectin, and report viral load results for fluvoxamine but not ivermectin. Presentation: falsely describing positive but not statistically significant effects as "no effect, what so ever" [Amrhein, odysee.com]. Prior statements: [odysee.com].

López-Medina et al.

An open letter, signed by >100 physicians, concluding this study is fatally flawed can be found at [jamaletter.com].

This is a phone survey based RCT with low risk patients, 200 ivermectin and 198 control, showing lower mortality, lower disease progression, lower treatment escalation, and faster resolution of symptoms with treatment, without reaching statistical significance. Authors find the results of this trial alone do not support the use of ivermectin. However the effects are all positive, especially for serious outcomes which are unable to reach statistical significance with the very small number of events in the low risk population.

With the low risk patient population, there is little room for improvement with an effective treatment - 59/57% (IVM/control) recovered within the first 2 days to either "no symptoms" or "not hospitalized and no limitation of activities"; 73/69% within 5 days. Less than 3% of all patients ever deteriorated.

The primary outcome was changed mid-trial, it was originally clinical deterioration, which is more meaningful, and shows greater benefit. The new outcome of resolution of symptoms includes "not hospitalized and no limitation of activities" as a negative outcome and is not very meaningful in terms of assessing how much treatment reduces serious outcomes. Using this measure could completely invalidate results - for example a treatment that eliminates all COVID-19 symptoms but has a temporary minor adverse event could be seen as worse.

Authors state that "preliminary reports of other randomized trials of ivermectin as treatment for COVID-19 with positive results have not yet been published in peer-reviewed journals", however there were 8 peer-reviewed RCTs with positive effects published prior to this paper (and 19 total peer-reviewed studies with positive effects).

Authors advised taking ivermectin on an empty stomach, reducing lung tissue concentration by ~2.5x [Guzzo].

76 patients were excluded due to control patients receiving ivermectin. However, there was a similar percentage of adverse events like diarrhea, nausea, and abdominal pain in both treatment and control groups. These are potential non-serious side effects of treatment and suggest that it is possible that many more control patients received some kind of treatment.

Ivermectin was widely used in the population and available OTC at the time of the study. The study protocol only excluded patients with previous ivermectin use within 5 days, however other trials often monitor effects 10+ days after the last dose [osf.io].

This study reportedly has an ethical issue whereby participants were told the study drug was "D11AX22" <https://trialsitenews.com/jama-ivermectin-study-deceived-participants-on-study-drug/>. The editor-in-chief of JAMA initially offered to help with this issue, but later indicated that "JAMA does not review consent forms", however the lead author reportedly confirmed the issue [[francesoir.fr](#), [trialsitenews.com](#), [trialsitenews.com](#) (B)].

The study protocol specifically allows "the use of other treatments outside of clinical trials". The paper provides no information on what other treatments were used, but other treatments were commonly used at the time. Additionally, the control group did about 5x better than anticipated for deterioration, also suggesting that the control patients used some kind of treatment. Patients that enroll in such a study may be more likely to learn about and use other treatments, especially since they do not know if they are receiving the study medication.

The study protocol was amended 4 times. Amendments 2-4 are provided but amendment 1 is missing. Amendment 2 increased the inclusion criteria to within 7 days of onset, including more later stage patients and reducing the expected effectiveness. The trial protocol lists "the duration of supplemental oxygen" as an outcome but the results for this outcome are missing.

Grants and/or personal fees, including in some cases during the conduct of the study, were provided by Sanofi Pasteur, GlaxoSmithKline, Janssen, Merck, and Gilead. For more details see [[trialsitenews.com](#) (C)].

For other confounding issues see [[osf.io](#) (B)] and additional issues can be found in the comments of the article [[jamanetwork.com](#)].

Gonzalez et al.

Another study reports results on a larger group of patients in the same hospital, showing ivermectin mortality RR 0.81 [0.53-1.24] [[Guzman](#)].

Questions have been raised about this study and the early termination of the study and discontinuation of treatments, because the hospital statistics show a dramatically lower (~75%) case fatality rate during the period of the study [[twitter.com](#)] (data from [[gob.mx](#)]).

Date	Cases	Deaths	CFR
3/2020	2	1	50%
4/2020	4	1	25%
5/2020	13	1	8%
6/2020	37	2	5%
7/2020	65	5	8%
8/2020	79	23	29%
9/2020	54	12	22%
10/2020	62	21	34%
11/2020	80	26	33%
12/2020	41	13	32%

Popp et al.

This is a very biased meta analysis designed to exclude almost all studies. Authors select a small subset of studies, with a majority of results based on only 1 or 2 studies, showing positive (non-statistically significant) results for 8 of 9 outcomes across a total of 13 studies. 5 outcomes are based on a single study, and 4 are based on 2 studies.

This analysis splits up studies in order to dilute the effects and avoid statistical significance. However, we can consider the probability of 8 of 9 positive effects occurring due to chance for an ineffective treatment, which is very unlikely (0.02 with an independence assumption).

The study is entirely retrospective in the current version. The protocol is dated April 20, 2021, and the most recent study included is from March 9, 2021. The protocol was modified after publication in order to include a close to null result (Gonzalez et al. "patients discharged without respiratory deterioration or death at 28 days"), so the current protocol is dated July 28, 2021.

Authors excluded many studies by requiring results at a specific time, for example mortality, ventilation, etc. required results at exactly 28 days. Authors excluded all prophylaxis studies by requiring results at exactly 14 days.

Studies comparing with other medications were excluded, however these studies confirm efficacy of ivermectin. The only case where they could overstate the efficacy of ivermectin is if the other medication was harmful. There is some evidence of this for excessive dosage/very late stage use, however that does not apply to any of the studies here.

Studies using combined treatment were excluded, even when it is known that the other components have minimal or no effect. 3 of 4 RCTs with combined treatment use doxycycline in addition [[sciencedirect.com](https://www.sciencedirect.com)]. Other studies were excluded by requiring PCR confirmation.

Authors are inconsistent regarding active comparators. They state that hydroxychloroquine "does not work", yet excluded trials comparing ivermectin to a drug they hold to be inactive. On the other hand, remdesivir was an acceptable comparator, although it is considered to be effective standard of care in some locations [osf.io (C)].

Authors fail to recognize that Risk of Bias (RoB) domains such as blinding are far less important for the objective outcome of mortality.

Bryant et al. note several other issues [osf.io (C)].

Cochrane was reputable in the past, but is now controlled by pharmaceutical interests. For example, see the news related to the expulsion of founder Dr. Gøtzsche and the associated mass resignation of board members in protest [blogs.bmj.com, bmj.com, en.x-mol.com]. For another example of bias see [ebm.bmj.com].

The BiRD group gave the following early comment: "Yesterday's Cochrane review surprisingly doesn't take a pragmatic approach comparing ivermectin versus no ivermectin, like in the majority of other existing reviews. It uses a granular approach similar to WHO's and the flawed Roman et al paper, splitting studies up and thereby diluting effects. Consequently, the uncertain conclusions add nothing to the evidence base. A further obfuscation of the evidence on ivermectin and an example of research waste. Funding conflicts of interests of the authors and of the journal concerned should be examined."

Roman et al.

This is a severely flawed meta analysis. An open letter signed by 40 physicians detailing errors and flaws, and requesting retraction, can be found at [\(trialsitenews.com \(D\)\)](https://trialsitenews.com). See also [\(bird-group.org\)](https://bird-group.org).

Authors cherry-pick to include only 4 studies reporting non-zero mortality and they initially claimed a mortality RR of 1.11 [0.16-7.65]. However, they reported incorrect values for Niaee et al., claiming an RR of 6.51 [2.18-19.45], when the correct RR for Niaee et al. is 0.18 [0.06-0.55]. After correction, their cherry-picked studies show >60% mortality reduction, however authors did not correct the conclusion.

Similarly, for viral clearance and NCT04392713, they report 20/41 treatment, 18/45 control, whereas the correct day 7 clearance numbers are 37/41 and 20/45 (sum of clearance @72hrs and @7 days), or 17/41 and 2/45 @72 hrs.

The duration of hospital stay for Niaee et al. is also incorrectly reported, showing a lower duration for the control group.

All of the errors are in one direction - incorrectly reporting lower than actual efficacy for ivermectin. Authors claim to include all RCTs excluding prophylaxis, however they only include 10 of the 24 non-prophylaxis RCTs (28 including prophylaxis at the time of publication). Authors actually reference meta analyses that do include the missing RCTs, so they should be aware of the missing RCTs.

The PubMed search strategy provided is syntactically incorrect. For additional errors, see [\(pubpeer.com\)](https://pubpeer.com). Also see [\(roundingtheearth.substack.com\)](https://roundingtheearth.substack.com).

The authors state that they have no conflicts of interest on medRxiv, however Dr. Pasupuleti's affiliation is Cello Health, whose website [\(cellohealth.com\)](https://cellohealth.com) notes that they provide services such as "brand and portfolio commercial strategy for biotech and pharma", and that their clients are "24 of the top 25 pharmaceutical companies".

Revisions

This paper is data driven, all graphs and numbers are dynamically generated. We will update the paper as new studies are released or with any corrections. Please submit updates and corrections at <https://ivmmeta.com/>.

11/26/2020: Initial revision.

12/2: We added [\[Ahmed\]](#).

12/7: We added [\[Chaccour\]](#).

12/11: We added [\[Soto-Becerra\]](#).

12/16: We added [\[Afsar\]](#).

12/17: We added [\[Alam\]](#).

12/26: We added [\[Carvallo \(B\), Vallejos\]](#).

12/27: We added the total number of authors and patients.

12/29: We added meta analysis excluding late treatment.

12/31: We added additional details about the studies in the appendix.

1/2/2021: We added dosage information and we added the number of patients to the forest plots.

1/5: We added direct links to the study details in the forest plots.

1/6: We added **[Babalola]**.

1/7: We added direct links to the study details in the chronological plots.

1/9: We added **[Ravikirti]**. Due to the much larger size of the control group in **[Bernigaud]**, we limited the size of the control group to be the same as the treatment group for calculation of the total number of patients.

1/10: We put all prophylaxis studies in a single group.

1/11: We added **[Chahla (B)]**.

1/12: We added **[Okumuş]**.

1/15: We added the effect measured for each study in the forest plots.

1/16: We moved the analysis with exclusions to the main text, and added additional commentary.

1/17: We added **[Bukhari]**.

1/19: We added **[Samaha, Shahbaznejad]**. **[Chaccour]** was updated to the journal version of the paper.

1/25: We updated **[Vallejos]** with the recently released results.

1/26: We updated **[Shouman]** with the journal version of the article.

2/2: We added **[Mohan]**.

2/5: We updated **[Bukhari]** to the preprint.

2/10: We added **[Lima-Morales]**.

2/11: We added more details on the analysis of prospective vs. retrospective studies.

2/12: We added **[Biber]**.

2/14: We added analysis restricted to COVID-19 case outcomes, and we added additional results in the abstract.

2/15: We added **[Behera (B)]**.

2/16: We updated **[Behera]** to the journal version of the paper.

2/17: We added **[Elalfy]**, and we added analysis restricted to viral clearance outcomes, and mortality results restricted to RCTs.

2/18: We updated **[Babalola]** to the journal version of the paper.

2/23: We added **[Gonzalez]**.

2/24: We added a comparison of the evidence base and WHO approval status for the use of ivermectin with scabies and COVID-19. We updated **[Okumuş]** with the Research Square preprint.

2/27: We added analysis restricted to peer reviewed studies.

3/2: We updated **[Vallejos]** with the latest results **[Vallejos (C)]**.

3/3: We updated the graphs to indicate the time period for the dosage column, now showing the dosage over one month for prophylaxis and over four days for other studies.

3/4: We added **[López-Medina]**, and we added more information in the abstract.

3/5: We added discussion of pooled effects (we show both pooled effects and individual outcome results).

3/6: We added **[Chowdhury]** and we identify studies that compare with another treatment.

3/10: We added **[Pott-Junior]**.

3/12: We added **[Bryant, Roy]**.

3/17: We added **[Nardelli]**.

3/25: We added **[Huvmek]**.

3/26: We added **[Tanioka]**.

3/28: We highlighted and added discussion for studies that use combined treatments.

3/30: We added **[Chahla]**.

3/31: We updated **[Chahla (B)]** to the preprint.

4/4: We added event counts to the forest plots.

4/5: We added **[Mourya]**.

4/7: We identified studies where minimal detail is currently available in the forest plots.

4/9: We corrected a duplicate entry for **[Bukhari]**.

4/10: We added **[Kishoria]**.

4/14: We added **[Seet]**.

4/16: We added **[Morgenstern]**.

4/18: We updated **[Morgenstern]** to the preprint.

4/25: We updated **[Biber]** to the latest results reported at the International Ivermectin for Covid Conference.

4/26: We added notes on heterogeneity.

4/27: We added analysis restricted to hospitalization results and a comparison with the evidence base used in the approval of other COVID-19 treatments.

4/28: We added the WHO meta analysis results for comparison.

4/30: We added analysis of the WHO meta analysis and updated **[Kory]** to the journal version.

5/4: We added **[Loue]**.

5/5: We previously limited the size of the control group in **[Bernigaud]** to be the same as the treatment group for calculation of the total number of patients. This is now also reflected and noted in the forest plots.

5/5: We updated **[Okumuş]** to the journal paper.

5/6: We updated discussion based on peer review including discussion of heterogeneity, exclusion based sensitivity analysis, and search criteria.

5/6: We added mechanical ventilation and ICU admission analysis.

5/6: We added a comparison of CDC recommendations.

5/6: We updated **[Chahla]** to the Research Square preprint.

5/7: We updated **[Shahbaznejad]** to the journal version, which includes additional outcomes not reported earlier.

5/8: We added **[Merino]**.

5/10: We added additional information in the abstract.

5/10: We added **[Faisal]**.

5/13: We updated **[Mahmud]** to the journal version.

5/15: We updated the discussion of the WHO analysis.

5/17: We added **[Szente Fonseca]**.

5/18: We added analysis of Merck's recommendation.

5/26: **[Samaha]** was updated to the journal version.

5/31: **[Biber]** was updated to the preprint.

6/2: We added **[Abd-El salam]**.

6/5: We added **[Ahsan]**.

6/7: We added **[Hariyanto]**.

6/15: We added **[Aref]**.

6/18: We added **[Krolewiecki]**.

6/19: **[Gonzalez]** was incorrectly included in the peer-reviewed analysis.

6/19: We updated **[Bryant]** to the journal version.

6/21: We added more information to the abstract.

7/2: We updated **[Niaee]** to the journal version.

7/3: We added **[Vallejos (B)]**.

7/6: We previously limited the size of the control group for **[Bernigaud]** when calculating the total number of patients, however this was confusing for many people that did not read the details. We now show the original counts and note the larger size of the control group in the text.

7/8: We updated **[Cadegiani]** to the journal version.

7/9: We added **[Hazan]**.

7/15: Elgazzar et al. was retracted and has been removed.

7/16: We updated **[Ravikirti]** with the journal version of the article.

7/20: We updated **[Hashim]** with the journal version of the article.

7/29: We added discussion in the responses section.

7/31: We added discussion in the responses section related to *in vitro* evidence and therapeutic concentrations.

8/2: We added analysis restricted to serious outcomes and analysis restricted to recovery, and we added discussion in the responses section.

8/3: We added discussion in the responses section.

8/4: We added discussion of the FDA recommendation.

8/5: We added **[Mondal]**.

8/6: We updated **[Behera (B)]** with the journal version of the article.

8/8: We updated discussion in the responses.

8/12: We added **[Elavarasi, Together Trial]**.

8/15: We updated discussion and made the abstract more concise.

8/16: We updated **[Together Trial]** with event counts.

8/26: We updated **[Mohan]** with the journal version of the article.

8/27: We updated **[Morgenstern (B)]** with the journal version of the article.

9/3: We updated discussion and excluded Carvallo et al. in the exclusion analysis.

9/6: We corrected **[Espitia-Hernandez]** to use the reported recovery time and added missing recovery and viral clearance results.

9/7: Discussion updates.

9/9: We added sensitivity analysis to compute the minimum number of studies that need to be excluded in order to avoid showing efficacy. Discussion updates.

9/14: FDA discussion updates.

9/15: Discussion updates.

9/17: We added study notes.

9/18: We added **[Buonfrate]**, and updated discussion of the Together Trial.

9/20: Discussion updates.

9/22: Discussion updates.

9/24: We added a graph of variants over time for the Together Trial discussion and corrected outcome discussion for Popp et al.

9/27: We added **[Mayer]**.

9/29: We corrected a display error causing a few points to be missing in Figure 3.

10/4: Merck discussion updates.

Appendix 1. Methods and Study Results

We performed ongoing searches of PubMed, medRxiv, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Collabovid, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19ivermectin.com, which regularly receives submissions of studies upon publication. Search terms were ivermectin and COVID-19 or SARS-CoV-2, or simply ivermectin. Automated searches are performed every hour with notifications of new matches. The broad search terms result in a large volume of new studies on a daily basis which are reviewed for inclusion. All studies regarding the use of ivermectin for COVID-19 that report a comparison with a control group are included in the main analysis. Sensitivity analysis is performed, excluding studies with critical issues, epidemiological studies, and studies with minimal available information. This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in calculations for that study. For example, if effects for mortality and cases are both reported, the effect for mortality is used, this may be different to the effect that a study focused on. If symptomatic results are reported at multiple times, we used the latest time, for example if mortality results are provided at 14 days and 28 days, the results at 28 days are used. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms were not used (the next most serious outcome is used — no studies were excluded). For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcome is considered more important than PCR testing status. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available (after most or all patients have recovered there is no room for an effective treatment to do better). If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO₂ is more important than cough. When results provide an odds ratio, we computed the relative risk when possible, or converted to a relative risk according to [Zhang]. Reported confidence intervals and *p*-values were used when available, using adjusted values when provided. If multiple types of adjustments are reported including propensity score matching (PSM), the PSM results are used. When needed, conversion between reported *p*-values and confidence intervals followed [Altman, Altman (B)], and Fisher's exact test was used to calculate *p*-values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the sum of the correction factors equal to 1 [Sweeting]. Results are all expressed with RR < 1.0 suggesting effectiveness. Most results are the relative risk of something negative. If studies report relative times, results are expressed as the ratio of the time for the ivermectin group versus the time for the control group. Calculations are done in Python (3.9.7) with scipy (1.7.1), pythonmeta (1.23), numpy (1.21.2), statsmodels (0.12.2), and plotly (5.3.1).

The forest plots are computed using PythonMeta [Deng] with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case). The forest plots show simplified dosages for comparison, these are the total dose in the first four days for treatment, and the monthly dose for prophylaxis, for a 70kg person. For full dosage details see below.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment, and treatment started within 5 days after the onset of symptoms, although a shorter time may be preferable. Antivirals are typically only considered effective when used within a shorter timeframe, for example 0-36 or 0-48 hours for oseltamivir, with longer delays not being effective [McLean, Treanor].

Note that the size of the control group in [Bernigaud] is significantly larger than the treatment group. We previously limited the size to be the same as that of the treatment group for calculation of the number of patients, however this was confusing to many people that did not read the details.

A summary of study results is below. Please submit updates and corrections at <https://ivmmeta.com/>.

Early treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in calculations, which may differ from the effect a paper focuses on.

<p>[Afsar], 12/15/2020, retrospective, Pakistan, South Asia, preprint, 6 authors, dosage 12mg days 1-6.</p>	<p>duration of fever, 98.0% lower, relative time 0.02, $p < 0.001$, treatment 0 of 37 (0.0%), control 7 of 53 (13.2%), adjusted per study, continuity correction due to zero event (with reciprocal of the contrasting arm), logistic regression.</p>
<p>[Ahmed], 12/2/2020, Double Blind Randomized Controlled Trial, Bangladesh, South Asia, peer-reviewed, mean age 42.0, 15 authors, dosage 12mg days 1-5, ivermectin + doxycycline group took only a single dose of ivermectin.</p>	<p>risk of unresolved symptoms, 85.0% lower, RR 0.15, $p = 0.09$, treatment 0 of 17 (0.0%), control 3 of 19 (15.8%), continuity correction due to zero event (with reciprocal of the contrasting arm), day 7 fever ivermectin.</p>
	<p>risk of unresolved symptoms, 62.7% lower, RR 0.37, $p = 0.35$, treatment 1 of 17 (5.9%), control 3 of 19 (15.8%), day 7 fever ivermectin + doxycycline.</p>
	<p>risk of no virological cure, 75.6% lower, RR 0.24, $p = 0.03$, treatment 11 of 22 (50.0%), control 20 of 23 (87.0%), adjusted per study, day 7 ivermectin.</p>
	<p>risk of no virological cure, 56.5% lower, RR 0.43, $p = 0.22$, treatment 16 of 23 (69.6%), control 20 of 23 (87.0%), adjusted per study, day 7 ivermectin + doxycycline.</p>
	<p>risk of no virological cure, 63.0% lower, RR 0.37, $p = 0.02$, treatment 5 of 22 (22.7%), control 14 of 23 (60.9%), adjusted per study, day 14 ivermectin.</p>
	<p>risk of no virological cure, 41.2% lower, RR 0.59, $p = 0.19$, treatment 9 of 23 (39.1%), control 14 of 23 (60.9%), adjusted per study, day 14 ivermectin + doxycycline.</p>
	<p>time to viral-, 23.6% lower, relative time 0.76, $p = 0.02$, treatment 22, control 23, ivermectin.</p>
	<p>time to viral-, 9.4% lower, relative time 0.91, $p = 0.27$, treatment 23, control 23, ivermectin + doxycycline.</p>
	<p>hospitalization time, 1.0% lower, relative time 0.99, ivermectin.</p>

	hospitalization time, 4.1% higher, relative time 1.04, ivermectin + doxycycline.
[Aref] , 6/15/2021, Randomized Controlled Trial, Egypt, Africa, peer-reviewed, 7 authors, dosage not specified.	relative duration of fever, 63.2% lower, relative time 0.37, $p < 0.001$, treatment 57, control 57.
	relative duration of dyspnea, 56.4% lower, relative time 0.44, $p < 0.001$, treatment 57, control 57.
	relative duration of anosmia, 68.8% lower, relative time 0.31, $p < 0.001$, treatment 57, control 57.
	relative duration of cough, 64.3% lower, relative time 0.36, $p < 0.001$, treatment 57, control 57.
	risk of no virological cure, 78.6% lower, RR 0.21, $p = 0.004$, treatment 3 of 57 (5.3%), control 14 of 57 (24.6%).
	time to viral-, 35.7% lower, relative time 0.64, $p < 0.001$, treatment 57, control 57.
[Babalola] , 1/6/2021, Double Blind Randomized Controlled Trial, Nigeria, Africa, peer-reviewed, baseline oxygen requirements 8.3%, 10 authors, dosage 12mg or 6mg q84h for two weeks, this trial compares with another treatment - results may be better when compared to placebo.	adjusted risk of viral+ at day 5, 63.9% lower, RR 0.36, $p = 0.11$, treatment 40, control 20, adjusted per study.
	relative ΔSpO_2 , 41.5% lower, RR 0.59, $p = 0.07$, treatment 38, control 18, figure 3.
	risk of no virological cure, 58.0% lower, RR 0.42, $p = 0.01$, treatment 20, control 20, 12mg - Cox proportional hazard model.
	risk of no virological cure, 40.5% lower, RR 0.60, $p = 0.12$, treatment 20, control 20, 6mg - Cox proportional hazard model.
	time to viral-, 49.2% lower, relative time 0.51, $p = 0.02$, treatment 20, control 20, 12mg.
	time to viral-, 34.4% lower, relative time 0.66, $p = 0.08$, treatment 20, control 20, 6mg.
[Biber] , 2/12/2021, Double Blind Randomized Controlled Trial, Israel, Middle East, preprint, 10 authors, dosage 12mg days 1-3, 15mg for patients $\geq 70\text{kg}$.	risk of hospitalization, 70.2% lower, RR 0.30, $p = 0.34$, treatment 1 of 47 (2.1%), control 3 of 42 (7.1%).
	risk of no virological cure, 44.8% lower, RR 0.55, $p = 0.04$, treatment 13 of 47 (27.7%), control 21 of

	42 (50.0%), adjusted per study, odds ratio converted to relative risk, multivariable logistic regression, day 6, Ct>30.
	risk of no virological cure, 70.2% lower, RR 0.30, $p = 0.14$, treatment 2 of 47 (4.3%), control 6 of 42 (14.3%), day 10, non-infectious samples (Ct>30 or non-viable culture).
	risk of no virological cure, 82.1% lower, RR 0.18, $p = 0.01$, treatment 2 of 47 (4.3%), control 10 of 42 (23.8%), day 8, non-infectious samples (Ct>30 or non-viable culture).
	risk of no virological cure, 75.6% lower, RR 0.24, $p = 0.02$, treatment 3 of 47 (6.4%), control 11 of 42 (26.2%), day 6, non-infectious samples (Ct>30 or non-viable culture).
	risk of no virological cure, 65.1% lower, RR 0.35, $p = 0.05$, treatment 4 of 28 (14.3%), control 9 of 22 (40.9%), day 4, non-infectious samples (Ct>30 or non-viable culture).
	risk of no virological cure, 51.9% lower, RR 0.48, $p = 0.08$, treatment 7 of 47 (14.9%), control 13 of 42 (31.0%), day 10, Ct>30.
	risk of no virological cure, 57.9% lower, RR 0.42, $p = 0.02$, treatment 8 of 47 (17.0%), control 17 of 42 (40.5%), day 8, Ct>30.
	risk of no virological cure, 44.7% lower, RR 0.55, $p = 0.05$, treatment 13 of 47 (27.7%), control 21 of 42 (50.0%), day 6, Ct>30.
	risk of no virological cure, 31.9% lower, RR 0.68, $p = 0.16$, treatment 13 of 28 (46.4%), control 15 of 22 (68.2%), day 4, Ct>30.
[Bukhari], 1/16/2021, Randomized Controlled Trial, Pakistan, South Asia, preprint, 10 authors, dosage 12mg single dose.	risk of no virological cure, 82.4% lower, RR 0.18, $p < 0.001$, treatment 4 of 41 (9.8%), control 25 of 45 (55.6%), day 7.
	risk of no virological cure, 38.7% lower, RR 0.61, $p < 0.001$, treatment 24 of 41 (58.5%), control 43 of 45 (95.6%), day 3.
[Buonfrate], 9/6/2021, Double Blind	risk of hospitalization, 600.0% higher, RR 7.00, $p =$

Randomized Controlled Trial, Italy, Europe, preprint, 18 authors, dosage 1200µg/kg days 1-5, arm B 600µg/kg, arm C 1200µg/kg.	0.30, treatment 4 of 58 (6.9%), control 0 of 29 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	relative change in viral load, RR 0.69, $p = 0.10$, treatment 30, control 29, day 7, arm C.
	relative change in viral load, RR 0.86, $p = 0.12$, treatment 28, control 29, day 7, arm B.
[Cadegiani] , 11/4/2020, prospective, Brazil, South America, peer-reviewed, 4 authors, dosage 200µg/kg days 1-3, this trial uses multiple treatments in the treatment arm (combined with AZ, nitazoxanide (82), HCQ (22), spironolactone (66), dutasteride (4)) - results of individual treatments may vary.	risk of death, 78.3% lower, RR 0.22, $p = 0.50$, treatment 0 of 110 (0.0%), control 2 of 137 (1.5%), continuity correction due to zero event (with reciprocal of the contrasting arm), control group 1.
	risk of mechanical ventilation, 94.2% lower, RR 0.06, $p = 0.005$, treatment 0 of 110 (0.0%), control 9 of 137 (6.6%), continuity correction due to zero event (with reciprocal of the contrasting arm), control group 1.
	risk of hospitalization, 98.0% lower, RR 0.02, $p < 0.001$, treatment 0 of 110 (0.0%), control 27 of 137 (19.7%), continuity correction due to zero event (with reciprocal of the contrasting arm), control group 1.
[Carvalho (C)] , 9/15/2020, prospective, Argentina, South America, peer-reviewed, mean age 55.7, 3 authors, dosage 36mg days 1, 8, dose varied depending on patient condition - mild 24mg, moderate 36mg, severe 48mg, this trial uses multiple treatments in the treatment arm (combined with dexamethasone, enoxaparin, and aspirin) - results of individual treatments may vary.	moderate/severe patients, 85.4% lower, RR 0.15, $p = 0.08$, treatment 1 of 32 (3.1%), control 3 of 14 (21.4%), the only treatment death was a patient already in the ICU before treatment.
[Chaccour] , 12/7/2020, Double Blind Randomized Controlled Trial, Spain, Europe, peer-reviewed, 23 authors, dosage 400µg/kg single dose.	risk of symptoms, 96.0% lower, RR 0.04, $p < 0.05$, treatment 12, control 12, logistic regression, chance of presenting any symptom, RR approximated with OR.
	viral load, 94.6% lower, relative load 0.05, $p < 0.01$, treatment 12, control 12, day 7 mid-recovery, average of gene E and gene N, data in supplementary appendix.
	risk of no virological cure, 8.0% lower, RR 0.92, $p =$

	1.00, treatment 12, control 12.
[Chahla] , 3/30/2021, Cluster Randomized Controlled Trial, Argentina, South America, preprint, 9 authors, dosage 24mg days 1, 8, 15, 22.	risk of no discharge, 86.9% lower, RR 0.13, $p = 0.004$, treatment 2 of 110 (1.8%), control 20 of 144 (13.9%), adjusted per study, odds ratio converted to relative risk, logistic regression.
[Chowdhury] , 7/14/2020, Randomized Controlled Trial, Bangladesh, South Asia, peer-reviewed, 6 authors, dosage 200µg/kg single dose, this trial compares with another treatment - results may be better when compared to placebo, this trial uses multiple treatments in the treatment arm (combined with doxycycline) - results of individual treatments may vary.	risk of hospitalization, 80.6% lower, RR 0.19, $p = 0.23$, treatment 0 of 60 (0.0%), control 2 of 56 (3.6%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	risk of no recovery, 46.4% lower, RR 0.54, $p < 0.001$, treatment 27 of 60 (45.0%), control 47 of 56 (83.9%), mid-recovery day 5.
	recovery time, 15.2% lower, relative time 0.85, $p = 0.07$, treatment 60, control 56.
	risk of no virological cure, 80.6% lower, RR 0.19, $p = 0.23$, treatment 0 of 60 (0.0%), control 2 of 56 (3.6%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	time to viral-, 4.3% lower, relative time 0.96, $p = 0.23$, treatment 60, control 56.
[Elalfy] , 2/16/2021, retrospective, Egypt, Africa, peer-reviewed, 15 authors, dosage 18mg days 1, 4, 7, 10, 13, <90kg 18mg, 90-120kg 24mg, >120kg 30mg, this trial uses multiple treatments in the treatment arm (combined with nitazoxanide, ribavirin, and zinc) - results of individual treatments may vary.	risk of no virological cure, 86.9% lower, RR 0.13, $p < 0.001$, treatment 7 of 62 (11.3%), control 44 of 51 (86.3%), day 15.
	risk of no virological cure, 58.1% lower, RR 0.42, $p < 0.001$, treatment 26 of 62 (41.9%), control 51 of 51 (100.0%), day 7.
[Espitia-Hernandez] , 8/15/2020, retrospective, Mexico, North America, peer-reviewed, mean age 45.1, 5 authors, dosage 6mg days 1-2, 8-9, this trial uses multiple treatments in the treatment arm (combined with azithromycin and cholecalciferol) - results of individual treatments may vary.	recovery time, 70.0% lower, relative time 0.30, $p < 0.001$, treatment 28, control 7.
	risk of viral+ at day 10, 97.2% lower, RR 0.03, $p < 0.001$, treatment 0 of 28 (0.0%), control 7 of 7 (100.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).
[Faisal] , 5/10/2021, Randomized Controlled Trial, Pakistan, South Asia, peer-reviewed, 3 authors, dosage 12mg days 1-5.	risk of no recovery, 68.4% lower, RR 0.32, $p = 0.005$, treatment 6 of 50 (12.0%), control 19 of 50 (38.0%), 6-8 days, mid-recovery.

	risk of no recovery, 27.3% lower, RR 0.73, $p = 0.11$, treatment 24 of 50 (48.0%), control 33 of 50 (66.0%), 3-5 days.
	risk of no recovery, 75.0% lower, RR 0.25, $p = 0.09$, treatment 2 of 50 (4.0%), control 8 of 50 (16.0%), 9-10 days.
[Krolewiecki] , 6/18/2021, Randomized Controlled Trial, Argentina, South America, peer-reviewed, 23 authors, dosage 600µg/kg days 1-5.	risk of mechanical ventilation, 151.9% higher, RR 2.52, $p = 1.00$, treatment 1 of 27 (3.7%), control 0 of 14 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	risk of disease progression, 3.7% higher, RR 1.04, $p = 1.00$, treatment 2 of 27 (7.4%), control 1 of 14 (7.1%).
	viral decay rate, 65.6% lower, RR 0.34, $p = 0.09$, treatment 20, control 14, relative mean viral decay rate (corrigendum table 2).
[Loue] , 4/17/2021, retrospective quasi-randomized (patient choice), France, Europe, peer-reviewed, 2 authors, dosage 200µg/kg single dose.	risk of death, 70.0% lower, RR 0.30, $p = 0.34$, treatment 1 of 10 (10.0%), control 5 of 15 (33.3%).
	risk of COVID-19 severe case, 55.0% lower, RR 0.45, $p = 0.11$, treatment 3 of 10 (30.0%), control 10 of 15 (66.7%).
[López-Medina] , 3/4/2021, Double Blind Randomized Controlled Trial, Colombia, South America, peer-reviewed, median age 37.0, 19 authors, dosage 300µg/kg days 1-5.	risk of death, 66.8% lower, RR 0.33, $p = 0.50$, treatment 0 of 200 (0.0%), control 1 of 198 (0.5%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	risk of escalation of care, 60.8% lower, RR 0.39, $p = 0.10$, treatment 4 of 200 (2.0%), control 10 of 198 (5.1%), odds ratio converted to relative risk.
	risk of escalation of care with post-hoc <12h exclusion, 34.3% lower, RR 0.66, $p = 0.51$, treatment 4 of 200 (2.0%), control 6 of 198 (3.0%), odds ratio converted to relative risk.
	risk of deterioration by ≥ 2 points on an 8-point scale, 43.1% lower, RR 0.57, $p = 0.35$, treatment 4 of 200 (2.0%), control 7 of 198 (3.5%), odds ratio converted to relative risk.
	risk of fever post randomization, 24.8% lower, RR 0.75, $p = 0.33$, treatment 16 of 200 (8.0%), control

	21 of 198 (10.6%), odds ratio converted to relative risk.
	risk of unresolved symptoms at day 21, 15.3% lower, RR 0.85, $p = 0.53$, treatment 36 of 200 (18.0%), control 42 of 198 (21.2%), odds ratio converted to relative risk, Cox proportional-hazard model.
	hazard ratio for lack of resolution of symptoms, 6.5% lower, RR 0.93, $p = 0.53$, treatment 200, control 198.
	relative median time to resolution of symptoms, 16.7% lower, relative time 0.83, treatment 200, control 198.
<p>[Mahmud], 10/9/2020, Double Blind Randomized Controlled Trial, Bangladesh, South Asia, peer-reviewed, 15 authors, dosage 12mg single dose, this trial uses multiple treatments in the treatment arm (combined with doxycycline) - results of individual treatments may vary.</p>	<p>risk of death, 85.7% lower, RR 0.14, $p = 0.25$, treatment 0 of 183 (0.0%), control 3 of 183 (1.6%), continuity correction due to zero event (with reciprocal of the contrasting arm).</p>
	risk of disease progression, 57.0% lower, RR 0.43, $p < 0.001$, treatment 16 of 183 (8.7%), control 32 of 180 (17.8%), adjusted per study, Cox regression.
	risk of no recovery, 94.0% lower, RR 0.06, $p < 0.001$, treatment 72 of 183 (39.3%), control 100 of 180 (55.6%), adjusted per study, day 7, Cox regression.
	risk of no recovery, 38.5% lower, RR 0.61, $p = 0.005$, treatment 40 of 183 (21.9%), control 64 of 180 (35.6%), day 11.
	risk of no recovery, 96.0% lower, RR 0.04, $p < 0.001$, treatment 42 of 183 (23.0%), control 67 of 180 (37.2%), adjusted per study, day 12, Cox regression.
	time to recovery, 27.0% lower, RR 0.73, $p = 0.003$, treatment 183, control 180, Cox regression.
	risk of no virological cure, 39.0% lower, RR 0.61, $p = 0.002$, treatment 14 of 183 (7.7%), control 36 of 180 (20.0%), adjusted per study, Cox regression.
<p>[Mayer], 9/23/2021, retrospective, Argentina, South America, preprint, 14</p>	<p>risk of death, 55.1% lower, RR 0.45, $p < 0.001$, treatment 3266, control 17966, adjusted per study,</p>

authors, dosage 540µg/kg days 1-5, mean prescribed dose.	odds ratio converted to relative risk, multiple logistic regression, Figure 3.
	risk of ICU admission, 65.9% lower, RR 0.34, $p < 0.001$, treatment 3266, control 17966, adjusted per study, odds ratio converted to relative risk, multiple logistic regression, Figure 3.
	risk of death, 27.6% lower, RR 0.72, $p = 0.03$, treatment 3266, control 17966, odds ratio converted to relative risk, unadjusted.
	risk of ICU admission, 26.0% lower, RR 0.74, $p = 0.13$, treatment 3266, control 17966, odds ratio converted to relative risk, unadjusted.
[Merino] , 5/3/2021, retrospective quasi-randomized (patients receiving kit), population-based cohort, Mexico, North America, preprint, 7 authors, dosage 6mg bid days 1-2.	risk of hospitalization, 74.4% lower, RR 0.26, $p < 0.001$, model 7, same time period, patients receiving kit.
	risk of hospitalization, 68.4% lower, RR 0.32, $p < 0.001$, model 1, different time periods, administrative rule.
[Mohan] , 2/2/2021, Double Blind Randomized Controlled Trial, India, South Asia, peer-reviewed, 27 authors, dosage 400µg/kg single dose, 200µg/kg also tested.	risk of no discharge at day 14, 62.5% lower, RR 0.38, $p = 0.27$, treatment 2 of 40 (5.0%), control 6 of 45 (13.3%), ivermectin 24mg.
	risk of no discharge at day 14, 43.8% lower, RR 0.56, $p = 0.49$, treatment 3 of 40 (7.5%), control 6 of 45 (13.3%), ivermectin 12mg.
	risk of clinical worsening, 32.5% lower, RR 0.68, $p = 0.72$, treatment 3 of 40 (7.5%), control 5 of 45 (11.1%), ivermectin 24mg.
	risk of clinical worsening, 55.0% lower, RR 0.45, $p = 0.44$, treatment 2 of 40 (5.0%), control 5 of 45 (11.1%), ivermectin 12mg.
	risk of no virological cure, 23.8% lower, RR 0.76, $p = 0.18$, treatment 21 of 40 (52.5%), control 31 of 45 (68.9%), ivermectin 24mg, day 5.
	risk of no virological cure, 5.6% lower, RR 0.94, $p = 0.82$, treatment 26 of 40 (65.0%), control 31 of 45 (68.9%), ivermectin 12mg, day 5.
	risk of no virological cure, 10.3% lower, RR 0.90, p

	<p>= 0.65, treatment 20 of 36 (55.6%), control 26 of 42 (61.9%), ivermectin 24mg, day 7.</p> <p>risk of no virological cure, 3.2% higher, RR 1.03, $p = 1.00$, treatment 23 of 36 (63.9%), control 26 of 42 (61.9%), ivermectin 12mg, day 7.</p>
[Mourya] , 4/1/2021, retrospective, India, South Asia, peer-reviewed, 5 authors, dosage 12mg days 1-7.	risk of no virological cure, 89.4% lower, RR 0.11, $p < 0.001$, treatment 5 of 50 (10.0%), control 47 of 50 (94.0%).
[Ravikirti] , 1/9/2021, Double Blind Randomized Controlled Trial, India, South Asia, peer-reviewed, 11 authors, dosage 12mg days 1, 2.	<p>risk of death, 88.7% lower, RR 0.11, $p = 0.12$, treatment 0 of 55 (0.0%), control 4 of 57 (7.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).</p> <p>risk of mechanical ventilation, 79.3% lower, RR 0.21, $p = 0.10$, treatment 1 of 55 (1.8%), control 5 of 57 (8.8%).</p> <p>risk of ICU admission, 13.6% lower, RR 0.86, $p = 0.80$, treatment 5 of 55 (9.1%), control 6 of 57 (10.5%).</p> <p>risk of no hospital discharge, 88.7% lower, RR 0.11, $p = 0.12$, treatment 0 of 55 (0.0%), control 4 of 57 (7.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).</p> <p>risk of no virological cure, 11.6% higher, RR 1.12, $p = 0.35$, treatment 42 of 55 (76.4%), control 39 of 57 (68.4%).</p>
[Roy] , 3/12/2021, retrospective, database analysis, India, South Asia, preprint, 5 authors, dosage not specified, this trial uses multiple treatments in the treatment arm (combined with doxycycline) - results of individual treatments may vary.	relative time to clinical response of wellbeing, 5.6% lower, relative time 0.94, $p = 0.87$, treatment 14, control 15.
[Samaha] , 1/16/2021, Randomized Controlled Trial, Lebanon, Middle East, peer-reviewed, 16 authors, dosage 12mg single dose, 45–64kg, 65–84kg, and >85kg patients received 9mg, 12mg, or 150µg/kg respectively.	<p>risk of hospitalization, 85.7% lower, RR 0.14, $p = 0.24$, treatment 0 of 50 (0.0%), control 3 of 50 (6.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).</p> <p>risk of fever at day 3, 90.9% lower, RR 0.09, $p = 0.004$, treatment 1 of 50 (2.0%), control 11 of 50 (22.0%).</p>

	relative improvement in Ct value at day 3, 68.3% lower, RR 0.32, $p < 0.001$, treatment 50, control 50.
[Szente Fonseca] , 10/31/2020, retrospective, Brazil, South America, peer-reviewed, mean age 50.6, 10 authors, dosage 12mg days 1-2.	risk of hospitalization, 13.9% higher, RR 1.14, $p = 0.45$, treatment 340, control 377, adjusted per study, odds ratio converted to relative risk, control prevalence approximated with overall prevalence.
[Together Trial] , 8/6/2021, Double Blind Randomized Controlled Trial, Brazil, South America, preprint, 1 author, dosage 400µg/kg days 1-3.	risk of death, 18.0% lower, RR 0.82, $p = 0.54$, treatment 18 of 677 (2.7%), control 22 of 678 (3.2%).
	extended ER observation or hospitalization, 9.0% lower, RR 0.91, $p = 0.51$, treatment 86 of 677 (12.7%), control 95 of 678 (14.0%).
[Vallejos (B)] , 7/2/2021, Double Blind Randomized Controlled Trial, Argentina, South America, peer-reviewed, 29 authors, dosage 12mg days 1-2, <80kg 12mg, 80-110kg 18mg, >110kg 24mg.	risk of death, 33.5% higher, RR 1.33, $p = 0.70$, treatment 4 of 250 (1.6%), control 3 of 251 (1.2%), odds ratio converted to relative risk.
	risk of mechanical ventilation, 33.5% higher, RR 1.33, $p = 0.70$, treatment 4 of 250 (1.6%), control 3 of 251 (1.2%), odds ratio converted to relative risk.
	risk of hospitalization, 33.0% lower, RR 0.67, $p = 0.23$, treatment 14 of 250 (5.6%), control 21 of 251 (8.4%), odds ratio converted to relative risk.
	risk of no virological cure, 5.0% higher, RR 1.05, $p = 0.55$, treatment 137 of 250 (54.8%), control 131 of 251 (52.2%), odds ratio converted to relative risk, day 3.
	risk of no virological cure, 26.8% higher, RR 1.27, $p = 0.29$, treatment 38 of 250 (15.2%), control 30 of 251 (12.0%), odds ratio converted to relative risk, day 12.

Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in calculations, which may differ from the effect a paper focuses on.

[Abd-Elsalam] , 6/2/2021, Randomized Controlled Trial, Egypt, Africa, peer-reviewed, 16 authors, dosage 12mg days 1-3.	risk of death, 25.0% lower, RR 0.75, $p = 0.70$, treatment 3 of 82 (3.7%), control 4 of 82 (4.9%), odds ratio converted to relative risk, logistic regression.
---	---

	<p>risk of mechanical ventilation, no change, RR 1.00, $p = 1.00$, treatment 3 of 82 (3.7%), control 3 of 82 (3.7%).</p> <p>hospitalization time, 19.6% lower, relative time 0.80, $p = 0.09$, treatment 82, control 82.</p>
[Ahsan] , 4/29/2021, retrospective, Pakistan, South Asia, peer-reviewed, 10 authors, dosage 150µg/kg days 1-2, 150-200µg/kg, this trial uses multiple treatments in the treatment arm (combined with doxycycline) - results of individual treatments may vary.	risk of death, 50.0% lower, RR 0.50, $p = 0.03$, treatment 17 of 110 (15.5%), control 17 of 55 (30.9%).
[Budhiraja] , 11/18/2020, retrospective, India, South Asia, preprint, 12 authors, dosage not specified.	risk of death, 99.1% lower, RR 0.009, $p = 0.04$, treatment 0 of 34 (0.0%), control 103 of 942 (10.9%), continuity correction due to zero event (with reciprocal of the contrasting arm), unadjusted.
[Camprubi] , 11/11/2020, retrospective, Spain, Europe, peer-reviewed, 9 authors, dosage 200µg/kg single dose.	risk of mechanical ventilation, 40.0% lower, RR 0.60, $p = 0.67$, treatment 3 of 13 (23.1%), control 5 of 13 (38.5%).
	risk of ICU admission, 33.3% lower, RR 0.67, $p = 1.00$, treatment 2 of 13 (15.4%), control 3 of 13 (23.1%), ICU at day 8.
	risk of no improvement at day 8, 33.3% higher, RR 1.33, $p = 1.00$, treatment 4 of 13 (30.8%), control 3 of 13 (23.1%).
	risk of no virological cure, 25.0% higher, RR 1.25, $p = 1.00$, treatment 5 of 13 (38.5%), control 4 of 13 (30.8%), tests done between days 3-5.
[Chachar] , 9/30/2020, Randomized Controlled Trial, India, South Asia, peer-reviewed, 6 authors, dosage 36mg, 12mg stat, 12mg after 12 hours, 12mg after 24 hours.	risk of no recovery at day 7, 10.0% lower, RR 0.90, $p = 0.50$, treatment 9 of 25 (36.0%), control 10 of 25 (40.0%).
[Elavarasi] , 8/12/2021, retrospective, India, South Asia, preprint, 26 authors, dosage not specified.	risk of death, 19.6% lower, RR 0.80, $p = 0.12$, treatment 48 of 283 (17.0%), control 311 of 1475 (21.1%), unadjusted.
[Gonzalez] , 2/23/2021, Double Blind Randomized Controlled Trial, Mexico,	risk of death, 14.4% lower, RR 0.86, $p = 1.00$, treatment 5 of 36 (13.9%), control 6 of 37 (16.2%).

North America, preprint, mean age 53.8, 13 authors, dosage 12mg single dose, 18mg for patients >80kg.	risk of respiratory deterioration or death, 8.6% lower, RR 0.91, $p = 1.00$, treatment 8 of 36 (22.2%), control 9 of 37 (24.3%).
	risk of no hospital discharge, 37.0% higher, RR 1.37, $p = 0.71$, treatment 4 of 36 (11.1%), control 3 of 37 (8.1%).
	hospitalization time, 20.0% higher, relative time 1.20, $p = 0.43$, treatment 36, control 37.
[Gorial] , 7/8/2020, retrospective, Iraq, Middle East, preprint, 9 authors, dosage 200µg/kg single dose.	risk of death, 71.0% lower, RR 0.29, $p = 1.00$, treatment 0 of 16 (0.0%), control 2 of 71 (2.8%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	hospitalization time, 42.0% lower, relative time 0.58, $p < 0.001$, treatment 16, control 71.
	risk of no recovery, 71.0% lower, RR 0.29, $p = 1.00$, treatment 0 of 16 (0.0%), control 2 of 71 (2.8%), continuity correction due to zero event (with reciprocal of the contrasting arm).
[Hashim] , 10/26/2020, Single Blind Randomized Controlled Trial, Iraq, Middle East, peer-reviewed, 7 authors, dosage 200µg/kg days 1-2, some patients received a third dose on day 8, this trial uses multiple treatments in the treatment arm (combined with doxycycline) - results of individual treatments may vary.	risk of death, 91.7% lower, RR 0.08, $p = 0.03$, treatment 0 of 59 (0.0%), control 6 of 70 (8.6%), continuity correction due to zero event (with reciprocal of the contrasting arm), excluding non-randomized critical patients.
	risk of death, 67.1% lower, RR 0.33, $p = 0.16$, treatment 2 of 70 (2.9%), control 6 of 70 (8.6%), odds ratio converted to relative risk, including critical patients that were always allocated to treatment.
	risk of disease progression, 83.1% lower, RR 0.17, $p = 0.07$, treatment 1 of 59 (1.7%), control 7 of 70 (10.0%), excluding non-randomized critical patients.
	risk of disease progression, 57.4% lower, RR 0.43, $p = 0.20$, treatment 3 of 70 (4.3%), control 7 of 70 (10.0%), odds ratio converted to relative risk, including critical patients that were always allocated to treatment.
	recovery time, 40.7% lower, relative time 0.59, $p <$

	0.001, treatment 70, control 70.
<p>[Hazan], 7/7/2021, retrospective, USA, North America, preprint, 7 authors, dosage 12mg days 1, 4, 8, this trial uses multiple treatments in the treatment arm (combined with doxycycline, zinc, vitamin D, vitamin C) - results of individual treatments may vary.</p>	<p>risk of death, 85.9% lower, RR 0.14, $p = 0.04$, continuity correction due to zero event (with reciprocal of the contrasting arm).</p>
	<p>risk of hospitalization, 93.3% lower, RR 0.07, $p = 0.001$, continuity correction due to zero event (with reciprocal of the contrasting arm).</p>
<p>[Huvemek], 3/25/2021, Double Blind Randomized Controlled Trial, Bulgaria, Europe, preprint, 1 author, dosage 400µg/kg days 1-3.</p>	<p>risk of no improvement, 31.6% lower, RR 0.68, $p = 0.28$, treatment 13 of 50 (26.0%), control 19 of 50 (38.0%), day 7, patients with improvement on WHO scale.</p>
	<p>risk of no improvement, 34.5% lower, RR 0.66, $p = 0.07$, treatment 19 of 50 (38.0%), control 29 of 50 (58.0%), day 4, patients with improvement on WHO scale.</p>
<p>[Khan], 9/24/2020, retrospective, Bangladesh, South Asia, preprint, median age 35.0, 8 authors, dosage 12mg single dose.</p>	<p>risk of death, 87.1% lower, RR 0.13, $p = 0.02$, treatment 1 of 115 (0.9%), control 9 of 133 (6.8%).</p>
	<p>risk of ICU admission, 89.5% lower, RR 0.11, $p = 0.007$, treatment 1 of 115 (0.9%), control 11 of 133 (8.3%).</p>
	<p>risk of disease progression, 83.5% lower, RR 0.17, $p < 0.001$, treatment 3 of 115 (2.6%), control 21 of 133 (15.8%).</p>
	<p>risk of no recovery, 87.1% lower, RR 0.13, $p = 0.02$, treatment 1 of 115 (0.9%), control 9 of 133 (6.8%).</p>
	<p>hospitalization time, 40.0% lower, relative time 0.60, $p < 0.001$, treatment 115, control 133.</p>
	<p>time to viral-, 73.3% lower, relative time 0.27, $p < 0.001$, treatment 115, control 133.</p>
<p>[Kishoria], 8/31/2020, Randomized Controlled Trial, India, South Asia, peer-reviewed, 7 authors, dosage 12mg single dose.</p>	<p>risk of no hospital discharge, 7.5% higher, RR 1.08, $p = 1.00$, treatment 11 of 19 (57.9%), control 7 of 13 (53.8%).</p>
	<p>risk of no virological cure, 7.5% higher, RR 1.08, $p = 1.00$, treatment 11 of 19 (57.9%), control 7 of 13 (53.8%), day 3.</p>
	<p>risk of no virological cure, 220.0% higher, RR 3.20,</p>

	<p>$p = 0.45$, treatment 1 of 5 (20.0%), control 0 of 6 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm), day 5.</p>
<p>[Lima-Morales], 2/10/2021, prospective, Mexico, North America, peer-reviewed, 9 authors, dosage 12mg single dose, this trial uses multiple treatments in the treatment arm (combined with azithromycin, montelukast, and aspirin) - results of individual treatments may vary.</p>	<p>risk of death, 77.7% lower, RR 0.22, $p < 0.001$, treatment 15 of 481 (3.1%), control 52 of 287 (18.1%), adjusted per study, odds ratio converted to relative risk, multivariate.</p>
	<p>risk of mechanical ventilation, 51.9% lower, RR 0.48, $p = 0.15$, treatment 8 of 434 (1.8%), control 11 of 287 (3.8%).</p>
	<p>risk of hospitalization, 67.4% lower, RR 0.33, $p < 0.001$, treatment 44 of 481 (9.1%), control 89 of 287 (31.0%), adjusted per study, odds ratio converted to relative risk, multivariate.</p>
	<p>risk of no recovery, 58.6% lower, RR 0.41, $p < 0.001$, treatment 75 of 481 (15.6%), control 118 of 287 (41.1%), adjusted per study, odds ratio converted to relative risk, recovery at day 14 after symptoms, multivariate.</p>
<p>[Niaee], 11/24/2020, Double Blind Randomized Controlled Trial, Iran, Middle East, peer-reviewed, mean age 56.0, 14 authors, dosage 400µg/kg single dose, dose varies in different groups.</p>	<p>risk of death, 81.8% lower, RR 0.18, $p = 0.001$, treatment 4 of 120 (3.3%), control 11 of 60 (18.3%), All IVM vs. all control.</p>
	<p>risk of death, 94.3% lower, RR 0.06, $p = 0.01$, treatment 0 of 30 (0.0%), control 11 of 60 (18.3%), continuity correction due to zero event (with reciprocal of the contrasting arm), IVM single dose 200mcg/kg vs. all control.</p>
	<p>risk of death, 45.5% lower, RR 0.55, $p = 0.37$, treatment 3 of 30 (10.0%), control 11 of 60 (18.3%), IVM three dose 200mcg/kg vs. all control.</p>
	<p>risk of death, 94.3% lower, RR 0.06, $p = 0.01$, treatment 0 of 30 (0.0%), control 11 of 60 (18.3%), continuity correction due to zero event (with reciprocal of the contrasting arm), IVM single dose 400mcg/kg vs. all control.</p>
	<p>risk of death, 81.8% lower, RR 0.18, $p = 0.06$, treatment 1 of 30 (3.3%), control 11 of 60 (18.3%), IVM three dose 400/200/200mcg/kg vs. all control.</p>

<p>[Okumuş], 1/12/2021, Double Blind Randomized Controlled Trial, Turkey, Europe, peer-reviewed, 15 authors, dosage 200µg/kg days 1-5, 36-50kg - 9mg, 51-65kg - 12mg, 66-79kg - 15mg, >80kg 200µg/kg.</p>	<p>risk of death, 33.3% lower, RR 0.67, $p = 0.55$, treatment 6 of 30 (20.0%), control 9 of 30 (30.0%).</p>
	<p>risk of no improvement at day 10, 42.9% lower, RR 0.57, $p = 0.18$, treatment 8 of 30 (26.7%), control 14 of 30 (46.7%).</p>
	<p>risk of no improvement at day 5, 15.8% lower, RR 0.84, $p = 0.60$, treatment 16 of 30 (53.3%), control 19 of 30 (63.3%).</p>
	<p>risk of no virological cure, 80.0% lower, RR 0.20, $p = 0.02$, treatment 2 of 16 (12.5%), control 5 of 8 (62.5%), day 10.</p>
<p>[Podder], 9/3/2020, Randomized Controlled Trial, Bangladesh, South Asia, peer-reviewed, 4 authors, dosage 200µg/kg single dose.</p>	<p>recovery time from enrollment, 16.1% lower, relative time 0.84, $p = 0.34$, treatment 32, control 30.</p>
<p>[Pott-Junior], 3/9/2021, Randomized Controlled Trial, Brazil, South America, peer-reviewed, 10 authors, dosage 200µg/kg single dose, dose varies in three arms 100, 200, 400µg/kg.</p>	<p>risk of mechanical ventilation, 85.2% lower, RR 0.15, $p = 0.25$, treatment 1 of 27 (3.7%), control 1 of 4 (25.0%).</p>
	<p>risk of ICU admission, 85.2% lower, RR 0.15, $p = 0.25$, treatment 1 of 27 (3.7%), control 1 of 4 (25.0%).</p>
	<p>relative improvement in Ct value, 0.8% lower, RR 0.99, $p = 1.00$, treatment 27, control 3.</p>
	<p>risk of no virological cure, 11.1% higher, RR 1.11, $p = 1.00$, treatment 10 of 27 (37.0%), control 1 of 3 (33.3%).</p>
	<p>time to viral-, 16.7% lower, relative time 0.83, treatment 27, control 3.</p>
<p>[Rajter], 10/13/2020, retrospective, propensity score matching, USA, North America, peer-reviewed, 6 authors, dosage 200µg/kg single dose.</p>	<p>risk of death, 46.0% lower, RR 0.54, $p = 0.04$, treatment 13 of 98 (13.3%), control 24 of 98 (24.5%), adjusted per study, odds ratio converted to relative risk, PSM.</p>
	<p>risk of death, 66.9% lower, RR 0.33, $p = 0.03$, treatment 26 of 173 (15.0%), control 27 of 107 (25.2%), adjusted per study, odds ratio converted to relative risk, multivariate.</p>
	<p>risk of mechanical ventilation, 63.6% lower, RR</p>

	0.36, $p = 0.10$, treatment 4 of 98 (4.1%), control 11 of 98 (11.2%), matched cohort excluding intubated at baseline.
[Shahbaznejad] , 1/19/2021, Double Blind Randomized Controlled Trial, Iran, Middle East, peer-reviewed, 8 authors, dosage 200µg/kg single dose.	risk of death, 197.1% higher, RR 2.97, $p = 1.00$, treatment 1 of 35 (2.9%), control 0 of 34 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm), patient died within 24 hours of admission.
	risk of mechanical ventilation, 94.3% higher, RR 1.94, $p = 1.00$, treatment 2 of 35 (5.7%), control 1 of 34 (2.9%).
	recovery time, 31.6% lower, relative time 0.68, $p = 0.05$, treatment 35, control 34, duration of dsypnea.
	recovery time, 19.2% lower, relative time 0.81, $p = 0.02$, treatment 35, control 34, duration of all symptoms.
	hospitalization time, 15.5% lower, relative time 0.85, $p = 0.02$, treatment 35, control 34.
[Soto-Becerra] , 10/8/2020, retrospective, database analysis, Peru, South America, preprint, median age 59.4, 4 authors, dosage 200µg/kg single dose.	risk of death, 17.1% lower, RR 0.83, $p = 0.01$, treatment 92 of 203 (45.3%), control 1438 of 2630 (54.7%), IVM vs. control day 43 (last day available) weighted KM from figure 3, per the pre-specified rules, the last available day mortality results have priority.
	risk of death, 39.0% higher, RR 1.39, $p = 0.16$, treatment 47 of 203 (23.2%), control 401 of 2630 (15.2%), adjusted per study, day 30, Table 2, IVM wHR.
[Spoorthi] , 11/14/2020, prospective, India, South Asia, peer-reviewed, 2 authors, dosage not specified, this trial uses multiple treatments in the treatment arm (combined with doxycycline) - results of individual treatments may vary.	recovery time, 21.1% lower, relative time 0.79, $p = 0.03$, treatment 50, control 50.
	hospitalization time, 15.5% lower, relative time 0.84, $p = 0.01$, treatment 50, control 50.

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in calculations, which may differ from the effect a paper focuses on.

[Alam] , 12/15/2020, prospective, Bangladesh, South Asia, peer-reviewed, 13 authors, dosage 12mg monthly.	risk of COVID-19 case, 90.6% lower, RR 0.09, $p < 0.001$, treatment 4 of 58 (6.9%), control 44 of 60 (73.3%).
[Behera (B)] , 2/15/2021, prospective, India, South Asia, peer-reviewed, 14 authors, dosage 300µg/kg days 1, 4.	risk of COVID-19 case, 83.0% lower, RR 0.17, $p < 0.001$, treatment 45 of 2199 (2.0%), control 133 of 1147 (11.6%), two doses.
	risk of COVID-19 case, 4.0% higher, RR 1.04, $p = 0.85$, treatment 23 of 186 (12.4%), control 133 of 1147 (11.6%), patients only receiving the first dose.
[Behera] , 11/3/2020, retrospective, India, South Asia, peer-reviewed, 13 authors, dosage 300µg/kg days 1, 4.	risk of COVID-19 case, 53.8% lower, RR 0.46, $p < 0.001$, treatment 41 of 117 (35.0%), control 145 of 255 (56.9%), adjusted per study, odds ratio converted to relative risk, model 2 2+ doses conditional logistic regression.
	risk of COVID-19 case, 44.5% lower, RR 0.56, $p < 0.001$, treatment 41 of 117 (35.0%), control 145 of 255 (56.9%), odds ratio converted to relative risk, matched pair analysis.
[Bernigaud] , 11/28/2020, retrospective, France, Europe, peer-reviewed, 12 authors, dosage 200µg/kg days 1, 8, 15, 400µg/kg days 1, 8, 15, two different dosages.	risk of death, 99.4% lower, RR 0.006, $p = 0.08$, treatment 0 of 69 (0.0%), control 150 of 3062 (4.9%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	risk of COVID-19 case, 55.1% lower, RR 0.45, $p = 0.01$, treatment 7 of 69 (10.1%), control 692 of 3062 (22.6%).
[Carvallo] , 11/17/2020, prospective, Argentina, South America, peer-reviewed, 4 authors, dosage 12mg weekly, this trial uses multiple treatments in the treatment arm (combined with iota-carrageenan) - results of individual treatments may vary.	risk of COVID-19 case, 99.9% lower, RR 0.001, $p < 0.001$, treatment 0 of 788 (0.0%), control 237 of 407 (58.2%), continuity correction due to zero event (with reciprocal of the contrasting arm).
[Carvallo (B)] , 10/19/2020, prospective, Argentina, South America, preprint, 1 author, dosage 1mg days 1-14, this trial uses multiple treatments in the	risk of COVID-19 case, 96.3% lower, RR 0.04, $p < 0.001$, treatment 0 of 131 (0.0%), control 11 of 98 (11.2%), continuity correction due to zero event (with reciprocal of the contrasting arm).

treatment arm (combined with iota-carrageenan) - results of individual treatments may vary.	
[Chahla (B)], 1/11/2021, Randomized Controlled Trial, Argentina, South America, peer-reviewed, 11 authors, dosage 12mg weekly, this trial uses multiple treatments in the treatment arm (combined with iota-carrageenan) - results of individual treatments may vary.	risk of moderate/severe case, 95.2% lower, RR 0.05, $p = 0.002$, treatment 0 of 117 (0.0%), control 10 of 117 (8.5%), continuity correction due to zero event (with reciprocal of the contrasting arm), moderate/severe COVID-19.
	risk of COVID-19 case, 84.0% lower, RR 0.16, $p = 0.001$, treatment 4 of 117 (3.4%), control 25 of 117 (21.4%), adjusted per study, odds ratio converted to relative risk, all cases.
[Hellwig], 11/28/2020, retrospective, ecological study, multiple countries, multiple regions, peer-reviewed, 2 authors, dosage 200µg/kg, dose varied, typically 150-200µg/kg.	risk of COVID-19 case, 78.0% lower, RR 0.22, $p < 0.02$, African countries, PCTI vs. no PCT, relative cases per capita.
	risk of COVID-19 case, 80.0% lower, RR 0.20, $p < 0.001$, worldwide, PCTI vs. no PCT, relative cases per capita.
[Mondal], 5/31/2021, retrospective, India, South Asia, peer-reviewed, 11 authors, dosage not specified.	risk of symptomatic case, 87.9% lower, RR 0.12, $p = 0.006$, treatment 128, control 1342, odds ratio converted to relative risk, multivariate logistic regression, control prevalence approximated with overall prevalence.
[Morgenstern], 4/16/2021, retrospective, propensity score matching, Dominican Republic, Caribbean, peer-reviewed, 16 authors, dosage 200µg/kg weekly.	risk of hospitalization, 80.0% lower, RR 0.20, $p = 0.50$, treatment 0 of 271 (0.0%), control 2 of 271 (0.7%), continuity correction due to zero event (with reciprocal of the contrasting arm), PSM.
	risk of COVID-19 case, 74.0% lower, RR 0.26, $p = 0.008$, treatment 5 of 271 (1.8%), control 18 of 271 (6.6%), adjusted per study, PSM, multivariate Cox regression.
[Seet], 4/14/2021, Cluster Randomized Controlled Trial, Singapore, Asia, peer-reviewed, 15 authors, dosage 12mg single dose, 200µg/kg, maximum 12mg, this trial compares with another treatment - results may be better when compared to placebo.	risk of symptomatic case, 49.8% lower, RR 0.50, $p = 0.01$, treatment 32 of 617 (5.2%), control 64 of 619 (10.3%).
	risk of COVID-19 case, 5.8% lower, RR 0.94, $p = 0.61$, treatment 398 of 617 (64.5%), control 433 of 619 (70.0%), adjusted per study, odds ratio converted to relative risk, model 6.
[Shouman], 8/28/2020, Randomized	risk of symptomatic case, 91.3% lower, RR 0.09, p

Controlled Trial, Egypt, Africa, peer-reviewed, 8 authors, dosage 18mg days 1, 3, dose varies depending on weight - 40-60kg: 15mg, 60-80kg: 18mg, >80kg: 24mg.	<p>< 0.001, treatment 15 of 203 (7.4%), control 59 of 101 (58.4%), adjusted per study, multivariate.</p> <p>risk of COVID-19 severe case, 92.9% lower, RR 0.07, $p = 0.002$, treatment 1 of 203 (0.5%), control 7 of 101 (6.9%), unadjusted.</p>
[Tanioka] , 3/26/2021, retrospective, ecological study, multiple countries, multiple regions, preprint, 3 authors, dosage 200µg/kg, dose varied, typically 150-200µg/kg.	<p>risk of death, 88.2% lower, RR 0.12, $p = 0.002$, relative mean mortality per million.</p>
[Vallejos] , 12/20/2020, retrospective, Argentina, South America, preprint, 1 author, dosage 12mg weekly.	<p>risk of COVID-19 case, 73.4% lower, RR 0.27, $p < 0.001$, treatment 13 of 389 (3.3%), control 61 of 486 (12.6%).</p>

References

1. **Abd-El salam** et al., Journal of Medical Virology, doi:10.1002/jmv.27122, *Clinical Study Evaluating the Efficacy of Ivermectin in COVID-19 Treatment: A Randomized Controlled Study*, <https://onlinelibrary.wiley.com/doi/10.1002/jmv.27122>.
2. **Adams**, B., Fierce Biotech, *Merck must do a new trial for faltering \$425M COVID-19 drug the U.S. government asked it to buy*, <https://www.fiercebiotech.com/biot..rug-u-s-government-asked-it-to-buy>.
3. **Afsar** et al., SSRN, *Ivermectin Use Associated with Reduced Duration of COVID-19 Febrile Illness in a Community Setting*, https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3734478.
4. **Ahmed** et al., International Journal of Infectious Diseases, doi:10.1016/j.ijid.2020.11.191, *A five day course of ivermectin for the treatment of COVID-19 may reduce the duration of illness*, <https://www.sciencedirect.com/science/article/pii/S1201971220325066>.
5. **Ahsan** et al., Cureus, doi:10.7759/cureus.14761, *Clinical Variants, Characteristics, and Outcomes Among COVID-19 Patients: A Case Series Analysis at a Tertiary Care Hospital in Karachi, Pakistan*, <https://www.cureus.com/articles/56..-care-hospital-in-karachi-pakistan>.
6. **Alam** et al., European Journal of Medical and Health Sciences, doi:10.24018/ejmed.2020.2.6.599, *Ivermectin as Pre-exposure Prophylaxis for COVID-19 among Healthcare Providers in a Selected Tertiary Hospital in Dhaka – An Observational Study*, <https://ejmed.org/index.php/ejmed/article/view/599>.
7. **Altman**, D., BMJ, doi:10.1136/bmj.d2304, *How to obtain the P value from a confidence interval*, <https://www.bmj.com/content/343/bmj.d2304>.
8. **Altman (B)** et al., BMJ, doi:10.1136/bmj.d2090, *How to obtain the confidence interval from a P value*, <https://www.bmj.com/content/343/bmj.d2090>.
9. **Amrhein** et al., Nature, 567:305-307, *Scientists rise up against statistical significance*, <https://www.nature.com/articles/d41586-019-00857-9>.

10. **Anglemyer** et al., Cochrane Database of Systematic Reviews 2014, Issue 4, doi:10.1002/14651858.MR000034.pub2, *Healthcare outcomes assessed with observational study designs compared with those assessed in randomized trials*, <https://www.cochranelibrary.com/cd..0.1002/14651858.MR000034.pub2/full>.
11. **Aref** et al., International Journal of Nanomedicine, doi:10.2147/IJN.S313093, *Clinical, Biochemical and Molecular Evaluations of Ivermectin Mucoadhesive Nanosuspension Nasal Spray in Reducing Upper Respiratory Symptoms of Mild COVID-19*, <https://www.dovepress.com/clinical..peer-reviewed-fulltext-article-IJN>.
12. **Arévalo** et al., Scientific Reports, doi:10.1038/s41598-021-86679-0 (preprint 11/2/20), *Ivermectin reduces in vivo coronavirus infection in a mouse experimental model*, <https://www.nature.com/articles/s41598-021-86679-0>.
13. **Babalola** et al., QJM: An International Journal of Medicine, doi:10.1093/qjmed/hcab035 (preprint 1/6), *Ivermectin shows clinical benefits in mild to moderate COVID19: A randomised controlled double-blind, dose-response study in Lagos*, <https://academic.oup.com/qjmed/adv../doi/10.1093/qjmed/hcab035/6143037>.
14. **Baqui** et al., The Lancet Global Health, doi:10.1016/S2214-109X(20)30285-0, *Ethnic and regional variations in hospital mortality from COVID-19 in Brazil: a cross-sectional observational study*, <https://www.sciencedirect.com/science/article/pii/S2214109X20302850>.
15. **Behera** et al., PLoS ONE, doi:10.1371/journal.pone.0247163 (preprint 11/3), *Role of ivermectin in the prevention of SARS-CoV-2 infection among healthcare workers in India: A matched case-control study*, <https://journals.plos.org/plosone/..le?id=10.1371/journal.pone.0247163>.
16. **Behera (B)** et al., Cureus 13:8, doi:10.7759/cureus.16897 (preprint 2/15/21), *Prophylactic Role of Ivermectin in Severe Acute Respiratory Syndrome Coronavirus 2 Infection Among Healthcare Workers*, <https://www.cureus.com/articles/64..infection-among-healthcare-workers>.
17. **Bello** et al., Journal of Biomolecular Structure and Dynamics, doi:10.1080/07391102.2021.1911857, *Elucidation of the inhibitory activity of ivermectin with host nuclear importin α and several SARS-CoV-2 targets*, <https://www.tandfonline.com/doi/full/10.1080/07391102.2021.1911857>.
18. **Bernigaud** et al., Annals of Dermatology and Venereology, doi:10.1016/j.annder.2020.09.231, *Ivermectin benefit: from scabies to COVID-19, an example of serendipity*, <https://www.sciencedirect.com/science/article/pii/S015196382030627X>.
19. **Bhattacharya** et al., Int. J. Scientific Research, doi:10.36106/ijsr/7232245, *Observational Study on Clinical Features, Treatment and Outcome of COVID 19 in a tertiary care Centre in India- a retrospective case series*, https://www.worldwidejournals.com/..ctober_2020_1614017661_0932284.pdf.
20. **Biber** et al., medRxiv, doi:10.1101/2021.05.31.21258081 (results 2/12/21), *Favorable outcome on viral load and culture viability using Ivermectin in early treatment of non-hospitalized patients with mild COVID-19, A double-blind, randomized placebo-controlled trial*, <https://www.medrxiv.org/content/10.1101/2021.05.31.21258081v1>.
21. **bird-group.org**, <https://bird-group.org/rebuttal-to-roman-et-al/>.
22. **blogs.bmj.com**, <https://blogs.bmj.com/bmj/2018/11/..ochrane-no-longer-a-collaboration/>.
23. **bmj.com**, <https://www.bmj.com/content/364/bmj.k5302>.
24. **Bray** et al., Antiviral Res., doi:10.1016/j.antiviral.2020.104805, *Ivermectin and COVID-19: A report in Antiviral Research, widespread interest, an FDA warning, two letters to the editor and the authors' responses*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7172803/>.
25. **Bryant** et al., American Journal of Therapeutics, doi:10.1097/MJT.0000000000001402 (preprint 3/11/21), *Ivermectin for Prevention and Treatment of COVID-19 Infection: A Systematic Review, Meta-analysis, and Trial Sequential Analysis to Inform Clinical Guidelines*,

https://journals.lww.com/american..Prevention_and_Treatment_of.7.aspx.

26. **Budhiraja** et al., medRxiv, doi:10.1101/2020.11.16.20232223, *Clinical Profile of First 1000 COVID-19 Cases Admitted at Tertiary Care Hospitals and the Correlates of their Mortality: An Indian Experience*, <https://www.medrxiv.org/content/10.1101/2020.11.16.20232223v1>.
27. **Bukhari** et al., medRxiv, doi:10.1101/2021.02.02.21250840 (results 1/16), *Efficacy of Ivermectin in COVID-19 Patients with Mild to Moderate Disease*, <https://www.medrxiv.org/content/10.1101/2021.02.02.21250840v1>.
28. **Buonfrate** et al., SSRN, doi:10.2139/ssrn.3918289, *High Dose Ivermectin for the Early Treatment of COVID-19 (COVIER Study): A Randomised, Double-Blind, Multicentre, Phase II, Dose-Finding, Proof of Concept Clinical Trial*, https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3918289.
29. **Cadegiani** et al., New Microbes and New Infections, doi:10.1016/j.nmni.2021.100915 (preprint 11/4/2020), *Early COVID-19 Therapy with azithromycin plus nitazoxanide, ivermectin or hydroxychloroquine in Outpatient Settings Significantly Improved COVID-19 outcomes compared to Known outcomes in untreated patients*, <https://www.sciencedirect.com/science/article/pii/S2052297521000792>.
30. **Caly** et al., Antiviral Research, doi:10.1016/j.antiviral.2020.104787, *The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro*, <https://www.sciencedirect.com/science/article/pii/S0166354220302011>.
31. **Camprubí** et al., PLoS ONE, 15:11, doi:10.1371/journal.pone.0242184, *Lack of efficacy of standard doses of ivermectin in severe COVID-19 patients*, <https://journals.plos.org/plosone/.le?id=10.1371/journal.pone.0242184>.
32. **Canga** et al., AAPS J., doi:10.1208/s12248-007-9000-9, *The Pharmacokinetics and Interactions of Ivermectin in Humans—A Mini-review*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2751445/>.
33. **Carvalho** et al., Journal of Biomedical Research and Clinical Investigation, doi:10.31546/2633-8653.1007, *Study of the Efficacy and Safety of Topical Ivermectin + Iota-Carrageenan in the Prophylaxis against COVID-19 in Health Personnel*, https://medicalpressopenaccess.com/upload/1605709669_1007.pdf.
34. **Carvalho (B)** et al., NCT04425850, *Usefulness of Topic Ivermectin and Carrageenan to Prevent Contagion of Covid 19 (IVERCAR)*, <https://clinicaltrials.gov/ct2/show/results/NCT04425850>.
35. **Carvalho (C)** et al., Journal of Clinical Trials, 11:459 (preprint 9/15/20), *Safety and Efficacy of the Combined Use of Ivermectin, Dexamethasone, Enoxaparin and Aspirina against COVID-19 the I.D.E.A. Protocol*, <https://www.longdom.org/open-access..vid19-the-idea-protocol-70290.html>.
36. **cellohealth.com**, <https://cellohealth.com/>.
37. **Chaccour** et al., EClinicalMedicine, doi:10.1016/j.eclinm.2020.100720 (preprint 12/7), *The effect of early treatment with ivermectin on viral load, symptoms and humoral response in patients with non-severe COVID-19: A pilot, double-blind, placebo-controlled, randomized clinical trial*, [https://www.thelancet.com/journals..PIIS2589-5370\(20\)30464-8/fulltext](https://www.thelancet.com/journals..PIIS2589-5370(20)30464-8/fulltext).
38. **Chaccour (B)** et al., Malar. J., doi:10.1186/s12936-017-1801-4, *Ivermectin to reduce malaria transmission I. Pharmacokinetic and pharmacodynamic considerations regarding efficacy and safety*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5402169/>.
39. **Chachar** et al., International Journal of Sciences, 9:31-35, doi:10.18483/ijSci.2378, *Effectiveness of Ivermectin in SARS-CoV-2/COVID-19 Patients*, <https://www.ijsciences.com/pub/article/2378>.
40. **Chahla** et al., Research Square, doi:10.21203/rs.3.rs-495945/v1 (original preprint 3/30), *Cluster Randomised Trials - Ivermectin Repurposing For COVID-19 Treatment Of Outpatients With Mild Disease In Primary Health Care Centers*, <https://www.researchsquare.com/article/rs-495945/v1>.

41. **Chahla (B)** et al., American Journal of Therapeutics, doi:10.1097/MJT.0000000000001433, *A randomized trial - intensive treatment based in ivermectin and iota-carrageenan as pre-exposure prophylaxis for COVID-19 in healthcare agents*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8415509/>.
42. **Chamie-Quintero** et al., OSF Preprints, *Ivermectin for COVID-19 in Peru: 14-fold reduction in nationwide excess deaths, $p=.002$ for effect by state, then 13-fold increase after ivermectin use restricted*, <https://osf.io/9egh4>.
43. **Chamie-Quintero (B)**, J., *The Latest Results of Ivermectin's Success in Treating Outbreaks of COVID-19*, <https://covid19criticalcare.com/iv..nalyse-on-covid19-and-ivermectin/>.
44. **Charilaou** et al., American Journal of Gastroenterology, doi:10.14309/00000434-201610001-01012, *Acetaminophen Toxicity: Trends in Hospitalization and Their Outcomes in United States from 2002-2011*, https://journals.lww.com/ajg/Fullt..rends_in_Hospitalization.1012.aspx.
45. **Choudhury** et al., Future Medicine, doi:10.2217/fvl-2020-0342, *Exploring the binding efficacy of ivermectin against the key proteins of SARS-CoV-2 pathogenesis: an in silico approach*, <https://www.futuremedicine.com/doi/10.2217/fvl-2020-0342>.
46. **Chowdhury** et al., Eurasian Journal of Medicine and Oncology, doi:10.14744/ejmo.2021.16263, *A Comparative Study on Ivermectin-Doxycycline and Hydroxychloroquine-Azithromycin Therapy on COVID-19 Patients*, <https://ejmo.org/10.14744/ejmo.2021.16263/>.
47. **Concato** et al., NEJM, 342:1887-1892, doi:10.1056/NEJM200006223422507, <https://www.nejm.org/doi/full/10.1056/nejm200006223422507>.
48. **Covid Analysis**, *Analysis of López-Medina et al.*, <https://c19ivermectin.com/lopezmedina.html>.
49. **COVID-NMA**, *COVID-NMA weekly update, May 14, 2021*, <https://web.archive.org/web/202105..58/https://www.covid-nma.com/news/>.
50. **de Melo** et al., EMBO Mol. Med., doi:10.15252/emmm.202114122 (preprint 11/22/20), *Attenuation of clinical and immunological outcomes during SARS-CoV-2 infection by ivermectin*, <https://www.embopress.org/doi/abs/10.15252/emmm.202114122>.
51. **Deaton** et al., Social Science & Medicine, 210, doi:10.1016/j.socscimed.2017.12.005, *Understanding and misunderstanding randomized controlled trials*, <https://www.sciencedirect.com/science/article/pii/S0277953617307359>.
52. **Deng, H.**, PyMeta, *Python module for meta-analysis*, <http://www.pymeta.com/>.
53. **Descotes, J.**, ImmunoSafe Consultance, *Medical Safety of Ivermectin*, https://www.medincell.com/wp-content..MDCL_safety_ivermectine-50321.pdf.
54. **DiNicolantonio** et al., Open Heart, doi:10.1136/openhrt-2020-001350, *Ivermectin may be a clinically useful anti-inflammatory agent for late-stage COVID-19*, <https://openheart.bmj.com/content/7/2/e001350>.
55. **DiNicolantonio (B)** et al., Open Heart, doi:10.1136/openhrt-2021-001655, *Anti-inflammatory activity of ivermectin in late-stage COVID-19 may reflect activation of systemic glycine receptors*, <https://openheart.bmj.com/content/8/1/e001655>.
56. **ebm.bmj.com**, <https://ebm.bmj.com/content/23/5/165>.
57. **Elalfy** et al., J. Med. Virol., doi:10.1002/jmv.26880, *Effect of a combination of Nitazoxanide, Ribavirin and Ivermectin plus zinc supplement (MANS.NRIZ study) on the clearance of mild COVID-1*, <https://onlinelibrary.wiley.com/doi/10.1002/jmv.26880>.

58. **Elavarasi** et al., medRxiv, doi:10.1101/2021.08.10.21261855, *Clinical features, demography and predictors of outcomes of SARS-CoV-2 infection in a tertiary care hospital in India - a cohort study*, <https://www.medrxiv.org/content/10.1101/2021.08.10.21261855v1>.
59. **en.x-mol.com**, <https://en.x-mol.com/paper/article/947631>.
60. **Errecalde** et al., Journal of Pharmaceutical Sciences, doi:10.1016/j.xphs.2021.01.017, *Safety and Pharmacokinetic Assessments of a Novel Ivermectin Nasal Spray Formulation in a Pig Model*, <https://www.sciencedirect.com/science/article/pii/S0022354921000320>.
61. **Espitia-Hernandez** et al., Biomedical Research, 31:5, *Effects of Ivermectin-azithromycin-cholecalciferol combined therapy on COVID-19 infected patients: A proof of concept study*, <https://www.biomedres.info/biomedinfo-proof-of-concept-study-14435.html>.
62. **Evans**, R., Healthcare Policy, 5:4, *Tough on Crime? Pfizer and the CIHR*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2875889/>.
63. **Eweas** et al., Frontiers in Microbiology, doi:10.3389/fmicb.2020.592908, *Molecular Docking Reveals Ivermectin and Remdesivir as Potential Repurposed Drugs Against SARS-CoV-2*, <https://www.frontiersin.org/articles/10.3389/fmicb.2020.592908/full>.
64. **Faisal** et al., The Professional Medical Journal, doi:10.29309/TPMJ/2021.28.05.5867, *Potential use of azithromycin alone and in combination with ivermectin in fighting against the symptoms of COVID-19*, <http://theprofesional.com/index.php/tpmj/article/view/5867>.
65. **Faria** et al., Science, doi:10.1126/science.abh2644, *Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil*, <https://www.science.org/lookup/doi/10.1126/science.abh2644>.
66. **Figuerola** et al., International Journal of General Medicine, doi:10.2147/IJGM.S328486 (preprint 4/15/2021), *Efficacy of a nasal spray containing Iota-Carrageenan in the prophylaxis of COVID-19 in hospital personnel dedicated to patients care with COVID-19 disease A pragmatic multicenter, randomized, double-blind, placebo-controlled trial (CARR-COV-02)*, <https://www.dovepress.com/efficacy.eer-reviewed-fulltext-article-IJGM>.
67. **Francés-Monerris** et al., ChemRxiv, doi:10.26434/chemrxiv.12782258.v1, *Has Ivermectin Virus-Directed Effects against SARS-CoV-2? Rationalizing the Action of a Potential Multitarget Antiviral Agent*, https://chemrxiv.org/articles/interp.target_Antiviral_Agent/12782258/1.
68. **francesoir.fr**, <https://www.francesoir.fr/societe...participants-letude-est-confirmer>.
69. **Galan** et al., Pathogens and Global Health, doi:10.1080/20477724.2021.1890887, *Phase 2 randomized study on chloroquine, hydroxychloroquine or ivermectin in hospitalized patients with severe manifestations of SARS-CoV-2 infection*, <https://www.tandfonline.com/doi/full/10.1080/20477724.2021.1890887>.
70. **Genetic Engineering and Biotechnology News**, *Merck to Seek FDA EUA for COVID-19 Pill after Positive Phase III Data*, <https://www.genengnews.com/news/merck-ill-after-positive-phase-iii-data/>.
71. **gob.mx**, <https://www.gob.mx/salud/documentos/datos-abiertos-152127>.
72. **Gonzalez** et al., medRxiv, doi:10.1101/2021.02.18.21252037, *Efficacy and safety of Ivermectin and Hydroxychloroquine in patients with severe COVID-19. A randomized controlled trial*, <https://www.medrxiv.org/content/10.1101/2021.02.18.21252037v1>.
73. **Gorial** et al., medRxiv, doi:10.1101/2020.07.07.20145979, *Effectiveness of Ivermectin as add-on Therapy in COVID-19 Management (Pilot Trial)*, <https://www.medrxiv.org/content/10.1101/2020.07.07.20145979v1>.
74. **Guzman** et al., medRxiv, doi:10.1101/2021.03.04.21252084, *Factors associated with increased mortality in critically ill COVID-19 patients in a Mexican public hospital: the other faces of health system oversaturation*, <https://www.medrxiv.org/content/10.1101/2021.03.04.21252084v1>.

75. **Guzzo** et al., J. Clinical Pharmacology, doi:10.1177/009127002237994, *Safety, Tolerability, and Pharmacokinetics of Escalating High Doses of Ivermectin in Healthy Adult Subjects*, <https://accp1.onlinelibrary.wiley...7/009127002237994?sid=nlm%3Apubmed>.
76. **Hariyanto** et al., Reviews In Medical Virology, doi:10.1002/rmv.2265, *Ivermectin and outcomes from Covid-19 pneumonia: A systematic review and meta-analysis of randomized clinical trial studies*, <https://onlinelibrary.wiley.com/doi/abs/10.1002/rmv.2265>.
77. **Hashim** et al., Iraqi Journal of Medical Science, 19:1, *Controlled randomized clinical trial on using Ivermectin with doxycycline for treating COVID-19 patients in Baghdad, Iraq*, <http://www.iraqijms.net/upload/pdf/iraqijms60db8b76d3b1e.pdf>.
78. **Hazan** et al., medRxiv, doi:10.1101/2021.07.06.21259924, *Effectiveness of Ivermectin-Based Multidrug Therapy in Severe Hypoxic Ambulatory COVID-19 Patients*, <https://www.medrxiv.org/content/10.1101/2021.07.06.21259924v1>.
79. **Heidary** et al., The Journal of Antibiotics, 73, 593–602, doi:10.1038/s41429-020-0336-z, *Ivermectin: a systematic review from antiviral effects to COVID-19 complementary regimen*, <https://www.nature.com/articles/s41429-020-0336-z>.
80. **Hellwig** et al., International Journal of Antimicrobial Agents, doi:10.1016/j.ijantimicag.2020.106248, *A COVID-19 Prophylaxis? Lower incidence associated with prophylactic administration of Ivermectin*, <https://www.sciencedirect.com/science/article/pii/S0924857920304684>.
81. **Huvemek** Press Release, *Kovid-19 - Huvemek® Phase 2 clinical trial*, <https://huvemec.bg/covid-19-huveme..linichno-izpitatie/za-isledvaneto/>.
82. **jamaletter.com**, <https://jamaletter.com/>.
83. **jamanetwork.com**, <https://jamanetwork.com/journals/jama/fullarticle/2777389>.
84. **Jans** et al., Cells 2020, 9:9, 2100, doi:10.3390/cells9092100, *Ivermectin as a Broad-Spectrum Host-Directed Antiviral: The Real Deal?*, <https://www.mdpi.com/2073-4409/9/9/2100>.
85. **Jeffreys** et al., bioRxiv, doi:10.1101/2020.12.23.424232, *Remdesivir-Ivermectin combination displays synergistic interaction with improved in vitro antiviral activity against SARS-CoV-2*, <https://www.biorxiv.org/content/10.1101/2020.12.23.424232v1>.
86. **Jerusalem Post**, *Israeli scientist says COVID-19 could be treated for under \$1/day*, <https://www.jpost.com/health-scienc..d-be-treated-for-under-1day-675612>.
87. **Kalfas** et al., medRxiv, doi:10.1101/2020.11.30.20236570, *The therapeutic potential of ivermectin for COVID-19: a systematic review of mechanisms and evidence*, <https://www.medrxiv.org/content/10.1101/2020.11.30.20236570v1>.
88. **Karita** et al., medRxiv, doi:10.1101/2021.08.27.21262754, *Trajectory of viral load in a prospective population-based cohort with incident SARS-CoV-2 G614 infection*, <https://www.medrxiv.org/content/10.1101/2021.08.27.21262754v1>.
89. **Khan** et al., Archivos de Bronconeumología, doi:10.1016/j.arbres.2020.08.007, *Ivermectin treatment may improve the prognosis of patients with COVID-19*, <https://www.archbronconeumol.org/e..ognosis-articulo-S030028962030288X>.
90. **Khan (B)**, T., PharmaShots, *Merck Signs ~\$1.2B Supply Agreement with US Government for Molnupiravir to Treat COVID-19*, <https://pharmashots.com/61076/merc..or-molnupiravir-to-treat-covid-19/>.

91. **Kishoria** et al., Paripex - Indian Journal of Research, doi:10.36106/paripex/4801859, *Ivermectin as adjuvant to hydroxychloroquine in patients resistant to standard treatment for SARS-CoV-2: results of an open-label randomized clinical study*, https://www.worldwidejournals.com/..August_2020_1597492974_4801859.pdf.
92. **Kory** et al., American Journal of Therapeutics, doi:10.1097/MJT.0000000000001377, *Review of the Emerging Evidence Demonstrating the Efficacy of Ivermectin in the Prophylaxis and Treatment of COVID-19*, https://journals.lww.com/american...Evidence_Demonstrating_the.4.aspx.
93. **Kory (B)**, P., *Commentary on medical journals*, <https://twitter.com/PierreKory/status/1434741303527579648>.
94. **Kory (C)**, P., *Dr. Pierre Kory Talks About Human Rights and The Big Science Disinformation*, <https://www.youtube.com/watch?v=3UTuT9TSRFQ&t=2890s>.
95. **Kory (D)**, P., *FLCCC Alliance Statement on the Irregular Actions of Public Health Agencies and the Widespread Disinformation Campaign Against Ivermectin*, <https://covid19criticalcare.com/wp..INFORMATION-CAMPAIGN-5.11.2021.pdf>.
96. **Krolewiecki** et al., EclinicalMedicine, doi:10.1016/j.eclinm.2021.100959, *Antiviral effect of high-dose ivermectin in adults with COVID-19: A proof-of-concept randomized trial*, <https://www.sciencedirect.com/science/article/pii/S258953702100239X>.
97. **Lawrie** et al., Preprint, *Ivermectin reduces the risk of death from COVID-19 – a rapid review and meta-analysis in support of the recommendation of the Front Line COVID-19 Critical Care Alliance*, <https://b3d2650e-e929-4448-a527-4e..b655bd21b1448ba6cf1f4c59f0d73d.pdf>.
98. **Lee** et al., Arch Intern Med., 2011, 171:1, 18-22, doi:10.1001/archinternmed.2010.482, *Analysis of Overall Level of Evidence Behind Infectious Diseases Society of America Practice Guidelines*, <https://jamanetwork.com/journals/j..nternalmedicine/fullarticle/226373>.
99. **Lehrer** et al., In Vivo, 34:5, 3023-3026, doi:10.21873/invivo.12134, *Ivermectin Docks to the SARS-CoV-2 Spike Receptor-binding Domain Attached to ACE2*, <http://iv.iarjournals.org/content/34/5/3023>.
100. **Li** et al., J. Cellular Physiology, doi:10.1002/jcp.30055, *Quantitative proteomics reveals a broad-spectrum antiviral property of ivermectin, benefiting for COVID-19 treatment*, <https://onlinelibrary.wiley.com/doi/10.1002/jcp.30055>.
101. **Lima-Morales**, *Effectiveness of a multidrug therapy consisting of ivermectin, azithromycin, montelukast and acetylsalicylic acid to prevent hospitalization and death among ambulatory COVID-19 cases in Tlaxcala, Mexico*, <https://www.sciencedirect.com/science/article/pii/S1201971221001004>.
102. **López-Medina** et al., JAMA, doi:10.1001/jama.2021.3071, *Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial*, <https://jamanetwork.com/journals/jama/fullarticle/2777389>.
103. **Loue** et al., J. Infectious Diseases and Epidemiology, doi:10.23937/2474-3658/1510202, *Ivermectin and COVID-19 in Care Home: Case Report*, <https://www.clinmedjournals.org/ar..idemiology-jide-7-202.php?jid=jide>.
104. **Madrid** et al., Heliyon, doi:10.1016/j.heliyon.2020.e05820, *Safety of oral administration of high doses of ivermectin by means of biocompatible polyelectrolytes formulation*, <https://www.sciencedirect.com/science/article/pii/S2405844020326633>.
105. **Mahmud** et al., Journal of International Medical Research, doi:10.5061/dryad.qjq2bvqf6 (preprint 10/9/20), *Ivermectin in combination with doxycycline for treating COVID-19 symptoms: a randomized trial*, <https://journals.sagepub.com/doi/10.1177/03000605211013550>.
106. **Mayer** et al., Zenodo, doi:10.5281/zenodo.5525362, *Safety and efficacy of a MEURI Program for the use of high dose ivermectin in COVID-19 patients*, <https://zenodo.org/record/5525362>.

107. **McLean** et al., *Open Forum Infect. Dis.* September 2015, 2:3, doi:10.1093/ofid/ofv100, *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4525010/>.
108. **Merck**, *Merck Statement on Ivermectin use During the COVID-19 Pandemic*, <https://www.merck.com/news/merck-s.-use-during-the-covid-19-pandemic/>.
109. **Merck (B)**, *Over 30 Years: The Mectizan® Donation Program*, <https://www.merck.com/stories/mectizan/>.
110. **Merck (C)** News Release, *Merck and Ridgeback's Investigational Oral Antiviral Molnupiravir Reduced the Risk of Hospitalization or Death by Approximately 50 Percent Compared to Placebo for Patients with Mild or Moderate COVID-19 in Positive Interim Analysis of Phase 3 Study*, <https://www.merck.com/news/merck-a.-for-patients-with-mild-or-moderat/>.
111. **Merino** et al., *SocArXiv Papers*, doi:10.31235/osf.io/r93g4, *Ivermectin and the odds of hospitalization due to COVID-19: evidence from a quasi-experimental analysis based on a public intervention in Mexico City*, <https://osf.io/preprints/socarxiv/r93g4/>.
112. **Miller**, K., *Dr. Fauci Says It's Fine to Take Vitamins C and D to Help Boost Your Immune System—Here's What to Know*, <https://www.health.com/nutrition/v.-pplements/dr-fauci-vitamin-c-and-d>.
113. **Mody** et al., *Communications Biology*, doi:10.1038/s42003-020-01577-x, *Identification of 3-chymotrypsin like protease (3CLPro) inhibitors as potential anti-SARS-CoV-2 agents*, <https://www.nature.com/articles/s42003-020-01577-x>.
114. **Mohan** et al., *Journal of Infection and Chemotherapy*, doi:10.1016/j.jiac.2021.08.021 (preprint 2/2/2021), *Single-dose oral ivermectin in mild and moderate COVID-19 (RIVET-COV): a single-centre randomized, placebo-controlled trial*, <https://www.sciencedirect.com/science/article/pii/S1341321X21002397>.
115. **Mondal** et al., *Journal of the Indian Medical Association*, 119:5, *Prevalence of COVID-19 Infection and Identification of Risk Factors among Asymptomatic Healthcare Workers: A Serosurvey Involving Multiple Hospitals in West Bengal*, https://onlinejima.com/read_journals.php?article=683.
116. **Morgenstern** et al., *Cureus*, doi:10.7759/cureus.17455 (preprint 4/16/2021), *Ivermectin as a SARS-CoV-2 Pre-Exposure Prophylaxis Method in Healthcare Workers: A Propensity Score-Matched Retrospective Cohort Study*, <https://www.cureus.com/articles/63..matched-retrospective-cohort-study>.
117. **Morgenstern (B)** et al., *J. Clinical Trials* (preprint 11/3), *The Use of Compassionate Ivermectin in the Management of Symptomatic Outpatients and Hospitalized Patients with Clinical Diagnosis of Covid-19 at the Centro Medico Bournigal and at the Centro Medico Punta Cana, Grupo Rescue, Dominican Republic, from May 1 to August 10, 2020*, <https://www.longdom.org/open-access..talized-patients-with-clinical.pdf>.
118. **Mountain Valley MD**, *Mountain Valley MD Receives Successful Results From BSL-4 COVID-19 Clearance Trial on Three Variants Tested With Ivectosol™*, <https://www.globenewswire.com/en/n..ariants-Tested-With-Ivectosol.html>.
119. **Mourya** et al., *Int. J. Health and Clinical Research*, *Comparative Analytical Study of Two Different Drug Regimens in Treatment of Covid 19 Positive Patients in Index Medical College Hospital and Research Center, Indore, India*, <https://ijhcr.com/index.php/ijhcr/article/view/1263>.
120. **Muñoz** et al., *PLoS Negl. Trop. Dis.*, doi:10.1371/journal.pntd.0006020, *Safety and pharmacokinetic profile of fixed-dose ivermectin with an innovative 18mg tablet in healthy adult volunteers*, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5773004/>.
121. **Nardelli** et al., *Signa Vitae*, doi:10.22514/sv.2021.043, *Crying wolf in time of Corona: the strange case of ivermectin and hydroxychloroquine. Is the fear of failure withholding potential life-saving treatment from clinical use?*, <https://www.signavitae.com/articles/10.22514/sv.2021.043>.

122. **Niaee** et al., Asian Pacific Journal of Tropical Medicine, doi:10.4103/1995-7645.318304 (preprint 11/24/20), *Ivermectin as an adjunct treatment for hospitalized adult COVID-19 patients: A randomized multi-center clinical trial*, <https://www.apjtm.org/article.asp?...66;epage=273;aulast=Shakhsi;type=0>.
123. **Nichol** et al., Injury, 2010, doi: 10.1016/j.injury.2010.03.033, *Challenging issues in randomised controlled trials*, [https://www.injuryjournal.com/article/S0020-1383\(10\)00233-0/fulltext](https://www.injuryjournal.com/article/S0020-1383(10)00233-0/fulltext).
124. **Nonaka** et al., International Journal of Infectious Diseases, doi:10.1016/j.ijid.2021.08.003, *SARS-CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil, February 2021*, <https://www.sciencedirect.com/science/article/pii/S1201971221006354>.
125. **O'Reilly**, D., *A Lifeline from Buenos Aires*, https://rescue.substack.com/p/li..ost&utm_medium=web&utm_source=copy.
126. **odysee.com**, <https://odysee.com/@FrontlineCovid..ents-defeating-the-delta-variant:a>.
127. **Okumuş** et al., BMC Infectious Diseases, doi:10.1186/s12879-021-06104-9 (preprint 1/12), *Evaluation of the Effectiveness and Safety of Adding Ivermectin to Treatment in Severe COVID-19 Patients*, <https://bmcinfectdis.biomedcentral..rticles/10.1186/s12879-021-06104-9>.
128. **Ontai** et al., medRxiv, doi:10.1101/2021.07.21.21260223, *Early multidrug treatment of SARS-CoV-2 (COVID-19) and decreased case fatality rates in Honduras*, <https://www.medrxiv.org/content/10.1101/2021.07.21.21260223v1>.
129. **Open Letter** by 170+ US Doctors, *JAMA Ivermectin Study Is Fatally Flawed*, <https://jamaletter.com/>.
130. **osf.io**, <https://osf.io/6m3ch/>.
131. **osf.io (B)**, <https://osf.io/bvznd>.
132. **osf.io (C)**, <https://osf.io/peqcj/>.
133. **otempo.com.br**, <https://www.otempo.com.br/interess..eocupa-por-risco-a-saude-1.2466432>.
134. **ourworldindata.org**, <https://ourworldindata.org/grapher/covid-variants-area?country=~BRA>.
135. **Pfeiffer**, M., *Analysis of the FDA's recommendation on ivermectin*, <https://twitter.com/marybethpf/status/1370182744718856193>.
136. **Pfizer**, *Pfizer Second Quarter 2021 Earnings Teleconference*, <https://s21.q4cdn.com/317678438/fi../Q2-2021-Earnings-Charts-FINAL.pdf>.
137. **Podder** et al., IMC J. Med. Science, 14:2, July 2020, *Outcome of ivermectin treated mild to moderate COVID-19 cases: a single-centre, open-label, randomised controlled study*, http://imcjms.com/registration/journal_abstract/353.
138. **Popp** et al., Cochrane Database of Systematic Reviews, doi:10.1002/14651858.CD015017.pub2, *Ivermectin for preventing and treating COVID-19*, <https://www.cochranelibrary.com/cd..0.1002/14651858.CD015017.pub2/full>.
139. **Pott-Junior** et al., Toxicology Reports, doi:10.1016/j.toxrep.2021.03.003, *Use of ivermectin in the treatment of Covid-19: a pilot trial*, <https://www.sciencedirect.com/science/article/pii/S2214750021000445>.
140. **pubpeer.com**, <https://pubpeer.com/publications/955418F3D4D39742CFFA8C1B023AA3>.
141. **Qureshi** et al., Journal of Biomolecular Structure and Dynamics, doi:10.1080/07391102.2021.1906750, *Mechanistic insights into the inhibitory activity of FDA approved ivermectin against SARS-CoV-2: old drug with new implications*, <https://www.tandfonline.com/doi/ab..02.2021.1906750?journalCode=tbsd20>.

142. **Rajter** et al., *Chest*, doi:10.1016/j.chest.2020.10.009, *Use of Ivermectin is Associated with Lower Mortality in Hospitalized Patients with COVID-19 (ICON study)*, <https://www.sciencedirect.com/science/article/pii/S0012369220348984>.
143. **Ravikirti** et al., *Journal of Pharmacy & Pharmaceutical Sciences*, doi:10.18433/jpps32105, *Ivermectin as a potential treatment for mild to moderate COVID-19: A double blind randomized placebo-controlled trial*, <https://journals.library.ualberta.../index.php/JPPS/article/view/32105>.
144. **reddit.com**, https://www.reddit.com/r/ivermecti..trial_was_not_randomized_possible/.
145. **Reuters**, *WHO joins Europe, Merck in recommending against ivermectin for COVID-19*, <https://news.trust.org/item/20210331135538-tajza/>.
146. **Rochweg** et al., *BMJ*, doi:10.1136/bmj.m2924, *Remdesivir for severe covid-19: a clinical practice guideline*, <https://www.bmj.com/content/370/bmj.m2924>.
147. **Roman** et al., *Clinical Infectious Diseases*, doi:10.1093/cid/ciab591 (preprint 5/25/21), *Ivermectin for the treatment of COVID-19: A systematic review and meta-analysis of randomized controlled trials*, <https://academic.oup.com/cid/advan..le/doi/10.1093/cid/ciab591/6310839>.
148. **roundingtheearth.substack.com**, <https://roundingtheearth.substack...ta-analytical-fixers-an-ivermectin>.
149. **Roy** et al., medRxiv, doi:10.1101/2021.03.08.21252883, *Outcome of Different Therapeutic Interventions in Mild COVID-19 Patients in a Single OPD Clinic of West Bengal: A Retrospective study*, <https://www.medrxiv.org/content/10.1101/2021.03.08.21252883v1>.
150. **Sabino** et al., *Lancet*, doi:10.1016/S0140-6736(21)00183-5, *Resurgence of COVID-19 in Manaus, Brazil, despite high seroprevalence*, [https://www.thelancet.com/article/S0140-6736\(21\)00183-5/fulltext](https://www.thelancet.com/article/S0140-6736(21)00183-5/fulltext).
151. **Saha** et al., *Structural Chemistry*, doi:10.1007/s11224-021-01776-0 (preprint 3/1), *The Binding mechanism of ivermectin and levosalbutamol with spike protein of SARS-CoV-2*, <https://www.researchsquare.com/article/rs-160254/v1>.
152. **Samaha** et al., *Viruses*, doi:10.3390/v13060989 (results 1/16), *Effects of a Single Dose of Ivermectin on Viral and Clinical Outcomes in Asymptomatic SARS-CoV-2 Infected Subjects: A Pilot Clinical Trial in Lebanon*, <https://www.mdpi.com/1999-4915/13/6/989/htm>.
153. **sciencedirect.com**, <https://www.sciencedirect.com/science/article/pii/S2213260021003106>.
154. **Seet** et al., *International Journal of Infectious Diseases*, doi:10.1016/j.ijid.2021.04.035, *Positive impact of oral hydroxychloroquine and povidone-iodine throat spray for COVID-19 prophylaxis: an open-label randomized trial*, [https://www.ijidonline.com/article/S1201-9712\(21\)00345-3/fulltext](https://www.ijidonline.com/article/S1201-9712(21)00345-3/fulltext).
155. **Shahbaznejad** et al., *Clinical Therapeutics*, doi:10.1016/j.clinthera.2021.04.007 (partial results available 1/19), *Effect of ivermectin on COVID-19: A multicenter double-blind randomized controlled clinical trial*, <https://www.sciencedirect.com/scie../article/abs/pii/S0149291821002010>.
156. **Shouman** et al., *Journal of Clinical and Diagnostic Research*, doi:10.7860/JCDR/2020/46795.0000, *Use of Ivermectin as a Potential Chemoprophylaxis for COVID-19 in Egypt: A Randomised Clinical Trial*, [https://www.jcdr.net/articles/PDF/..Sh\)_PF1\(SY_OM\)_PFA_\(OM\)_PN\(KM\).pdf](https://www.jcdr.net/articles/PDF/..Sh)_PF1(SY_OM)_PFA_(OM)_PN(KM).pdf).
157. **Soto-Becerra** et al., medRxiv, doi:10.1101/2020.10.06.20208066, *Real-World Effectiveness of hydroxychloroquine, azithromycin, and ivermectin among hospitalized COVID-19 patients: Results of a target trial emulation using observational data from a nationwide Healthcare System in Peru*, <https://www.medrxiv.org/content/10.1101/2020.10.06.20208066v1>.
158. **Spoorthi** et al., *IAIM*, 2020, 7:10, 177-182, *Utility of Ivermectin and Doxycycline combination for the treatment of SARSCoV-2*, http://iaimjournal.com/wp-content/..oads/2020/10/iaim_2020_0710_23.pdf.

159. **Surnar** et al., ACS Pharmacol. Transl. Sci., doi:10.1021/acsptsci.0c00179, *Clinically Approved Antiviral Drug in an Orally Administrable Nanoparticle for COVID-19*, <https://pubs.acs.org/doi/abs/10.1021/acsptsci.0c00179>.
160. **Sweeting** et al., Statistics in Medicine, doi:10.1002/sim.1761, *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, <https://onlinelibrary.wiley.com/doi/10.1002/sim.1761>.
161. **Szente Fonseca** et al., Travel Medicine and Infectious Disease, doi:10.1016/j.tmaid.2020.101906, *Risk of Hospitalization for Covid-19 Outpatients Treated with Various Drug Regimens in Brazil: Comparative Analysis*, <https://www.sciencedirect.com/scie..article/abs/pii/S1477893920304026>.
162. **Tanioka** et al., medRxiv, doi:10.1101/2021.03.26.21254377, *Why COVID-19 is not so spread in Africa: How does Ivermectin affect it?*, <https://www.medrxiv.org/content/10.1101/2021.03.26.21254377v1>.
163. **Together Trial**, *Early Treatment of COVID-19 with Repurposed Therapies: The TOGETHER Adaptive Platform Trial*, <https://rethinkingclinicaltrials.o..tform-trial-edward-mills-phd-frcp/>.
164. **Treanor** et al., JAMA, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016, *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, <https://jamanetwork.com/journals/jama/fullarticle/192425>.
165. **TrialSiteNews**, *Pfizer CEO All but Acknowledges Vaccines Won't be Enough – Here Comes the Antivirals & Billions More*, <https://trialsitenews.com/pfizer-c..omes-the-antivirals-billions-more/>.
166. **trialsitenews.com**, <https://trialsitenews.com/jama-una..tudy-consent-form-used-ivermectin/>.
167. **trialsitenews.com (B)**, <https://trialsitenews.com/jama-ive..articipants-is-publicly-confirmed/>.
168. **trialsitenews.com (C)**, <https://trialsitenews.com/is-there..g-the-trial-site-during-the-study/>.
169. **trialsitenews.com (D)**, <https://trialsitenews.com/statemen..iseases-acceptance-of-roman-et-al/>.
170. **twitter.com**, <https://twitter.com/jjchamie/status/1438631284163715074>.
171. **Udofia** et al., Network Modeling Analysis in Health Informatics and Bioinformatics, doi:10.1007/s13721-021-00299-2, *In silico studies of selected multi-drug targeting against 3CLpro and nsp12 RNA-dependent RNA-polymerase proteins of SARS-CoV-2 and SARS-CoV*, <https://link.springer.com/article/10.1007/s13721-021-00299-2>.
172. **Vallejos** et al., Preliminary Results, *Ivermectina en agentes de salud e IVERCOR COVID19*, <https://twitter.com/Covid19Crusher/status/1365420061859717124>.
173. **Vallejos (B)** et al., BMC Infectious Diseases, doi:10.1186/s12879-021-06348-5, *Ivermectin to prevent hospitalizations in patients with COVID-19 (IVERCOR-COVID19) a randomized, double-blind, placebo-controlled trial*, <https://bmcinfectdis.biomedcentral..rticles/10.1186/s12879-021-06348-5>.
174. **Vallejos (C)** et al., *Ivermectina en agentes de salud e ivercor COVID-19: resultados al 18 de feb 2021*, <https://twitter.com/Covid19Crusher/status/1365420061859717124>.
175. **Wagstaff** et al., Ivermectin Global Summit, *In vitro investigations of ivermectin as an antiviral agent*, <https://vimeo.com/554860553#t=1h32m0s>.
176. **Wehbe** et al., Front. Immunol., doi:10.3389/fimmu.2021.663586, *Repurposing Ivermectin for COVID-19: Molecular Aspects and Therapeutic Possibilities*, <https://www.frontiersin.org/articles/10.3389/fimmu.2021.663586/full>.
177. **WHO**, *Therapeutics and COVID-19: Living Guideline 31 March 2021*, <https://apps.who.int/iris/bitstrea..9-nCoV-therapeutics-2021.1-eng.pdf>.
178. **Wikipedia**, *Molnupiravir*, <https://en.wikipedia.org/wiki/Molnupiravir>.

179. **Yagisawa** et al., The Japanese Journal of Antibiotics, 74-1, Mar 2021, *Global trends in clinical studies of ivermectin in COVID-19*, http://jja-contents.wdc-jp.com/pdf/JJA74/74-1-open/74-1_44-95.pdf.
180. **Yesilbag** et al., Virus Research, doi:10.1016/j.virusres.2021.198384, *Ivermectin also inhibits the replication of bovine respiratory viruses (BRSV, BPIV-3, BoHV-1, BCoV and BVDV) in vitro*, <https://www.sciencedirect.com/science/article/pii/S0168170221000915>.
181. **Yim**, P., TrialSiteNews, *Systemic unreported protocol violations in key ivermectin study*, <https://trialsitenews.com/systemic-violations-in-key-ivermectin-study/>.
182. **Zaidi** et al., The Journal of Antibiotics, doi:10.1038/s41429-021-00430-5, *The mechanisms of action of Ivermectin against SARS-CoV-2: An evidence-based clinical review article*, <https://www.nature.com/articles/s41429-021-00430-5>.
183. **Zatloukal** et al., *News report on In Vitro results from the research institute of Prof. Zatloukal*, <https://www.servustv.com/videos/aa-27juub3a91w11/>.
184. **Zavascki** et al., Research Square, doi:10.21203/rs.3.rs-910467/v1, *Advanced ventilatory support and mortality in hospitalized patients with COVID-19 caused by Gamma (P.1) variant of concern compared to other lineages: cohort study at a reference center in Brazil*, <https://www.researchsquare.com/article/rs-910467/v1>.
185. **Zhang** et al., JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690, *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, <https://jamanetwork.com/journals/jama/fullarticle/188182>.